

Observation of the $B^+ \rightarrow J\psi\eta'K^+$ decay



The LHCb collaboration

E-mail: Ivan.Belyaev@cern.ch

ABSTRACT: The $B^+ \rightarrow J\psi\eta'K^+$ decay is observed for the first time using proton-proton collision data collected by the LHCb experiment at centre-of-mass energies of 7, 8, and 13 TeV, corresponding to a total integrated luminosity of 9 fb^{-1} . The branching fraction of this decay is measured relative to the known branching fraction of the $B^+ \rightarrow \psi(2S)K^+$ decay and found to be

$$\frac{\mathcal{B}(B^+ \rightarrow J\psi\eta'K^+)}{\mathcal{B}(B^+ \rightarrow \psi(2S)K^+)} = (4.91 \pm 0.47 \pm 0.29 \pm 0.07) \times 10^{-2},$$

where the first uncertainty is statistical, the second is systematic and the third is related to external branching fractions. A first look at the $J/\psi\eta'$ mass distribution is performed and no signal of intermediate resonances is observed.

KEYWORDS: B Physics, Branching fraction, Charm Physics, Hadron-Hadron Scattering

ARXIV EPRINT: [2303.09443](https://arxiv.org/abs/2303.09443)

Contents

1	Introduction	1
2	Detector and simulation	2
3	Event selection	3
4	Signal yield determination	5
5	Normalisation channel	7
6	Branching fraction ratio computation	8
7	Systematic uncertainties	9
8	Results and summary	10
	The LHCb collaboration	20

1 Introduction

In the last twenty years a plethora of new hadron states have been discovered in decays of beauty hadrons to charmonium, including the enigmatic $\chi_{c1}(3872)$ state [1], numerous pentaquark states in $J/\psi p$ [2–6] and $J/\psi \Lambda$ [7] systems as well as tetraquarks in the $\psi(2S)\pi^+$ [8–12], $J/\psi\phi$ [13–17], $\eta_c(1S)\pi^-$ [18], $J/\psi\pi^+$ [19], $J/\psi K^+$ [16] and $J/\psi K_S^0$ [17] systems. Transitions among charmonium or charmonium-like states have been studied in beauty-hadron decays, including transitions with emission of one photon [20–23], two pions [24–27], ϕ [13–17], ω [27, 28] and η [29] mesons. The transitions with emission of an η meson have also been studied in $e^+e^- \rightarrow J/\psi\eta$ processes [30, 31]. In general, studies of various hadronic transitions in the charmonium and charmonium-like sectors can shed light onto the internal structure of these particles, which is largely unknown for newly discovered hadronic states [32–35].

Transitions with an emission of an η' meson in the charmonium and charmonium-like systems have not yet been observed [32, 36, 37]. Since the η' meson may have a glueball contribution [38–55], processes involving this particle are of particular interest [56, 57]. The $B^+ \rightarrow J/\psi\eta'K^+$ decay¹ is a good candidate to explore the $J/\psi\eta'$ system in detail, offering the opportunity to search for possible intermediate resonances. The decay itself has never been observed and an upper limit on its branching fraction of

$$\mathcal{B}(B^+ \rightarrow J/\psi\eta'K^+) < 8.8 \times 10^{-5} \text{ (90 \% CL)},$$

was set by the Belle collaboration [58].

¹Inclusion of charge-conjugate states is implied throughout the paper, unless otherwise stated.

This paper reports the observation of the $B^+ \rightarrow J/\psi \eta' K^+$ decay using proton-proton (pp) collision data collected by the LHCb experiment at centre-of-mass energies of 7, 8, and 13 TeV, corresponding to a total integrated luminosity of 9 fb^{-1} . The measurement of its branching fraction normalised to the well-known branching fraction of the $B^+ \rightarrow \psi(2S) K^+$ decay [37],

$$\mathcal{R} \equiv \frac{\mathcal{B}(B^+ \rightarrow J/\psi \eta' K^+)}{\mathcal{B}(B^+ \rightarrow \psi(2S) K^+)} , \quad (1.1)$$

is performed using the $\eta' \rightarrow \rho^0 \gamma$ decay. The observation of the signal is confirmed using the $\eta' \rightarrow \eta \pi^+ \pi^-$ decay mode, which is also used as a cross-check.

2 Detector and simulation

The LHCb detector [59, 60] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region [61], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes [62, 63] placed downstream of the magnet. The tracking system provides a measurement of the momentum, p , of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at 200 GeV/ c . The minimum distance of a track to a primary pp collision vertex (PV), the impact parameter is measured with a resolution of $(15 + 29/p_T) \mu\text{m}$, where p_T is the component of the momentum transverse to the beam, in GeV/ c . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors [64]. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [65].

The online event selection is performed by a trigger [66, 67], which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction. At the hardware trigger stage, events are required to have a muon track with high transverse momentum or dimuon candidates in which the product of the p_T of the muons has a high value. In the software trigger, two oppositely charged muons are required to form a good-quality vertex that is significantly displaced from every PV, with a dimuon mass exceeding $2.7 \text{ GeV}/c^2$.

Simulated events are used to describe signal shapes and to compute the efficiencies needed to determine the branching fraction ratio. In the simulation, pp collisions are generated using PYTHIA [68] with a specific LHCb configuration [69]. Decays of unstable particles are described by EVTGEN [70], in which final-state radiation is generated using PHOTOS [71]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [72] as described in ref. [74]. The p_T and rapidity (y) spectra of the B^+ mesons in simulation are corrected to match distributions in data. The correction factors are calculated by comparing the observed p_T and y

spectra for a high-purity data sample of reconstructed $B^+ \rightarrow J/\psi K^+$ decays with the corresponding simulated samples. In the simulation, the $B^+ \rightarrow J/\psi \eta' K^+$ decays are generated as phase-space decays and corrected using a gradient boosted decision tree reweighting algorithm [75] to reproduce the $J/\psi \eta'$ and $\eta' K^+$ mass spectra observed in data. To describe accurately the variables used for kaon identification, the corresponding quantities in simulation are resampled according to values obtained from calibration data samples of $D^{*+} \rightarrow (D^0 \rightarrow K^- \pi^+) \pi^+$ decays [76]. The procedure accounts for correlations between the variables associated with a particular track, as well as the dependence of the kaon identification response on the track's p_T and η and the multiplicity of tracks in the event. To account for imperfections in the simulation of charged-particle reconstruction, the track reconstruction efficiency is corrected using a sample of $J/\psi \rightarrow \mu^+ \mu^-$ decays in data [77]. Samples of the $B^+ \rightarrow J/\psi K^{*+}$ decays with $K^{*+} \rightarrow K^+ (\pi^0 \rightarrow \gamma \gamma)$ are used to correct the photon reconstruction efficiency in simulation [78–81].

3 Event selection

The $B^+ \rightarrow J/\psi \eta' K^+$ candidates are reconstructed with η' decays to either $(\rho^0 \rightarrow \pi^+ \pi^-) \gamma$ or $(\eta \rightarrow \gamma \gamma) \pi^+ \pi^-$ final states. The difficulty of reconstructing photons in the $\eta \rightarrow \gamma \gamma$ decay leads to a sample with fewer events. The $B^+ \rightarrow \psi(2S) K^+$ normalisation decay is reconstructed using the $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ decay. In both signal and normalisation channels, the J/ψ meson is reconstructed in its decay to two muons. As explained in detail below, an initial loose selection is applied for both signal and normalisation channels. Subsequently, for the $B^+ \rightarrow J/\psi \eta' K^+$ candidates, where the background level is large, a multivariate estimator is used to select higher purity subset of candidates. The normalisation channel has a high purity after the initial selection, therefore no further selection steps are applied.

To reduce systematic uncertainties, the initial selection criteria for both signal and normalisation channels are kept the same whenever possible. The selection criteria are chosen to be similar to those used in previous LHCb studies [20, 21, 23, 29, 53, 78, 79]. The muon, pion and kaon candidates are identified by combining information from the Cherenkov detectors, calorimeters and muon detectors [82] associated with the reconstructed tracks. To reduce the combinatorial background, only tracks that are inconsistent with originating from any reconstructed PV in the event are considered. The transverse momentum of the muon candidates is required to be greater than 500 MeV/ c and their momenta must exceed 6 GeV/ c . Pairs of oppositely-charged muons consistent with originating from a common vertex are combined to form $J/\psi \rightarrow \mu^+ \mu^-$ candidates. The reconstructed mass of the muon pair is required to be between 3.056 and 3.136 GeV/ c^2 .

Tracks that are consistent with the pion or kaon hypotheses are required to have transverse momentum greater than 200 MeV/ c . Photons are reconstructed from clusters in the electromagnetic calorimeter, with transverse energy above 350 MeV. The clusters must not be associated with reconstructed tracks [83, 84]. Photon identification is based on the combined information from electromagnetic and hadronic calorimeters, scintillation pad, preshower detectors and the tracking system.

For the reconstruction of the $\eta' \rightarrow \eta \pi^+ \pi^-$ candidates, two photons are first combined to form an η candidate. The diphoton mass is restricted to lie within $\pm 60 \text{ MeV}/c^2$ around the known mass of the η meson [37]. Each η candidate is then combined with two oppositely-charged pions to form an η' candidate. The mass of the combination is required to lie within $\pm 45 \text{ MeV}/c^2$ around the known mass of the η' meson [37]. For an η' candidate reconstructed in the $\eta' \rightarrow \rho^0 \gamma$ decay mode, the ρ^0 candidate is formed from two oppositely-charged pions. The mass of this candidate is restricted to lie between 500 and 900 MeV/c^2 . This asymmetric region around the known mass of the ρ^0 meson [37] takes into account the shift of the ρ^0 line shape, due to the electric-dipole nature of the $\eta' \rightarrow \rho^0 \gamma$ transition [85–90]. A photon is combined with the ρ^0 candidate in order to form the η' candidate, whose mass is required to lie within $\pm 30 \text{ MeV}/c^2$ of the known η' mass [37].

Each selected J/ψ candidate is combined with a kaon track and either an η' candidate or two oppositely-charged pions to form a B^+ candidate decaying into the signal or normalisation modes, respectively. For the $B^+ \rightarrow \psi(2S)K^+$ candidates the $J/\psi \pi^+ \pi^-$ mass is required to be between 3.66 and 3.71 GeV/c^2 . To improve the B^+ meson mass resolution a kinematic fit [91] is performed, which constrains the masses of the J/ψ , η' and η candidates to their known values [37], and the B^+ candidates to originate from its associated PV. The decay time of the B^+ candidates is required to be greater than $100 \mu\text{m}/c$ to suppress the large combinatorial background from tracks created in a PV.

Further selection of the $B^+ \rightarrow J/\psi \eta' K^+$ decays is based on a multivariate estimator, in the following referred to as the multi-layer perceptron (MLP) classifier. The classifier is based on an artificial neural network algorithm [92, 93], configured with a cross-entropy cost estimator [94]. It reduces the combinatorial background to a low level while retaining a high signal efficiency. Two MLP classifiers are trained separately for the two different η' meson decay modes. The list of variables used for classifiers includes the χ^2 of the kinematic fit; transverse momenta of the η' , kaon and pion candidates; pseudorapidities of pion and kaon candidates; transverse momentum of the photon from the $\eta' \rightarrow \rho^0 \gamma$ decay or minimal transverse momentum of photons from the $\eta' \rightarrow (\eta \rightarrow \gamma \gamma) \pi^+ \pi^-$ decay; decay time of the B^+ candidate; variable related to the quality of kaon identification [64, 82] and, for the $\eta' \rightarrow \rho^0 \gamma$ decay, cosine of the angle between momenta of the π^+ and η' candidates in the rest frame of the ρ^0 candidate. The classifiers are trained using simulated samples of $B^+ \rightarrow J/\psi \eta' K^+$ decays as signal proxy, while the $B^+ \rightarrow J/\psi \eta' K^+$ candidates from data with mass above 5.35 GeV/c^2 are used to represent the background. The $B^+ \rightarrow J/\psi (\eta' \rightarrow \rho^0 \gamma) K^+$ candidates with the $J/\psi \pi^+ \pi^- K^+$ mass consistent with the known mass of the B^+ meson are vetoed to avoid contamination from the $B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$ decays with a random photon added.

The requirement on each of the MLP classifiers is chosen to maximise the figure-of-merit defined as $S/\sqrt{B+S}$, where S represents the expected signal yield, and B is the expected background yield within a $\pm 15 \text{ MeV}/c^2$ mass window centred around the known mass of the B^+ meson and corresponding to approximately three times the mass resolution on both sides of the peak. The background yield is calculated from fits to data, as described in section 4, while the expected signal yield is estimated as $S = \varepsilon S_0$, where ε is the efficiency of the requirement on the response of the MLP classifier determined from simulation,

and S_0 is the signal yield obtained from the fit to the data, with a loose requirement applied on the response of the MLP classifier.² The mass distributions for the selected $B^+ \rightarrow J/\psi \eta' K^+$ candidates are shown in figure 1, where clear signal peaks corresponding to B^+ decays are seen in data for both η' decay modes.

4 Signal yield determination

The signal yields are determined using an extended unbinned maximum-likelihood fit to the $J/\psi \eta' K^+$ mass distributions with a two-component function. The signal component for both cases is modelled by a modified Gaussian function that combines a Gaussian core with power-law tails on both sides of the distribution [95, 96]. The background component is parameterised by a second-order positive polynomial function [97] for the $\eta' \rightarrow \rho^0 \gamma$ decay mode and by an exponential function in case of the $\eta' \rightarrow \eta \pi^+ \pi^-$ decay mode. The parameters of the detector resolution function are taken from simulation, and the width of the Gaussian function is further corrected by a scale factor, s_{B^+} , that accounts for a small discrepancy between data and simulation [15, 26, 29, 98, 99]. To account for the uncertainty in the tail parameters and resolution, the fit is performed simultaneously for data and simulated samples, sharing the same tail parameters, and allowing the correction factor s_{B^+} to vary. The resulting fit functions are overlaid with data distributions in figure 1 and the signal yields are found to be

$$N_{B^+ \rightarrow J/\psi \eta' K^+} \Big|_{\eta' \rightarrow \rho^0 \gamma} = (1.11 \pm 0.11) \times 10^3, \quad (4.1a)$$

$$N_{B^+ \rightarrow J/\psi \eta' K^+} \Big|_{\eta' \rightarrow \eta \pi^+ \pi^-} = (0.228 \pm 0.028) \times 10^3, \quad (4.1b)$$

where the uncertainties are statistical only. The resolution correction factors s_{B^+} are found to be 1.08 ± 0.12 and 1.03 ± 0.16 for the $\eta' \rightarrow \rho^0 \gamma$ and $\eta' \rightarrow \eta \pi^+ \pi^-$ samples, respectively. In both cases, the statistical significance of the $B^+ \rightarrow J/\psi \eta' K^+$ signal is calculated using Wilks' theorem [100] and found to exceed 17 and 12 standard deviations for the $\eta' \rightarrow \rho^0 \gamma$ and $\eta' \rightarrow \eta \pi^+ \pi^-$ samples, respectively. However, as the signal yield is much lower for the η' meson decays to the $\eta \pi^+ \pi^-$ final state, all subsequent studies are performed using only the $\eta' \rightarrow \rho^0 \gamma$ decay mode.

The background-subtracted $J/\psi \eta'$, $\eta' K^+$, and $J/\psi K^+$ mass spectra from the $B^+ \rightarrow J/\psi (\eta' \rightarrow \rho^0 \gamma) K^+$ decays are shown in figures 2(a-c), where the *sPlot* technique [101] based on the fit results is used for background subtraction. The $J/\psi \eta'$, $\eta' K^+$, and $J/\psi K^+$ masses are calculated using a kinematic fit with J/ψ , η' and B^+ mass constraints and a PV constraint applied [91]. While for the $J/\psi K^+$ mass the distribution largely agrees with the shape expected from the phase-space model, for the low-mass region of the $\eta' K^+$ mass spectrum and the high-mass region of the $J/\psi \eta'$ mass spectrum a striking difference from the phase-space model is observed. These differences are potentially due to contributions from decays via intermediate heavy excited strange mesons, such as $K_0^*(1430)^+$, $K_2^*(1430)^+$ or $K^*(1680)^+$ mesons, decaying into the $\eta' K^+$ final state. The decays of the B^+ mesons into

²The optimal requirement is found to be largely independent on the choice of the normalisation point.




Figure 1. Mass distributions for selected $B^+ \rightarrow J/\psi \eta' K^+$ candidates with η' decays to (left) $\rho^0 \gamma$ and (right) $\eta \pi^+ \pi^-$ final states. The resulting fit functions are overlaid with data distributions.




Figure 2. Normalised background-subtracted (a) $\eta' K^+$, (b) $J/\psi \eta'$, (c) $J/\psi K^+$ mass spectra and (d) two-dimensional mass distribution of $\eta' K^+$ vs $J/\psi \eta'$ from the $B^+ \rightarrow J/\psi \eta' K^+$ decays. Superimposed curves are the expectations from a phase-space model.

a J/ψ meson and heavy excited strange mesons have been studied in refs. [13, 14, 16]. The decays via intermediate excited kaons also contribute to the higher mass region of the $J/\psi \eta'$ mass spectrum, as shown in figure 2(d). The $J/\psi \eta'$ mass region below $4.7 \text{ GeV}/c^2$ is explicitly inspected for possible contributions from decays via excited charmonium or charmoni-

um-like states into the $J/\psi\eta'$ final state. Fits to the background-subtracted $J/\psi\eta'$ mass distribution are performed in individual mass windows, corresponding to the well-established $\psi(4160)$, $\psi(4230)$, $\psi(4360)$, $\psi(4415)$ and $\psi(4430)$ resonances [37]. For each fit, the resonance shape is parameterised with a relativistic Breit-Wigner function convoluted with a mass resolution function. The non-resonant contribution is modelled by a first order positive polynomial function. The known masses and widths of the resonances [37] are introduced in the fits as Gaussian constraints on the corresponding parameters. The resolution function is modelled by the modified Gaussian function with parameters obtained using simulation as a function of the $J/\psi\eta'$ mass. No statistically significant signals are observed for the $B^+ \rightarrow J/\psi\eta'K^+$ decays via intermediate resonances, listed above. To probe the contribution of the resonances with higher masses, more advanced fit techniques accounting for complicated background shape and the distortion of the signal Breit-Wigner shape are required.

For the determination of the resonant structure of the $B^+ \rightarrow J/\psi\eta'K^+$ decay, a full amplitude analysis, similar to those used in refs. [14, 16], is required. Large signal yields and low background levels are important prerequisites for such analysis. The relatively large level of combinatorial background for the $B^+ \rightarrow J/\psi\eta'K^+$ signal decays with the η' meson reconstructed via the $\eta' \rightarrow \rho^0\gamma$ decay mode, makes it difficult to carry out an amplitude analysis. With a larger data sample, expected from future data-taking periods, it will be possible to perform the full amplitude analysis using the $B^+ \rightarrow J/\psi\eta'K^+$ signal decays with the η' meson reconstructed via the $\eta' \rightarrow \eta\pi^+\pi^-$ decay mode, where the combinatorial background is smaller.

5 Normalisation channel

The $J/\psi\pi^+\pi^-K^+$ mass distribution for selected $B^+ \rightarrow (\psi(2S) \rightarrow J/\psi\pi^+\pi^-)K^+$ candidates with $J/\psi\pi^+\pi^-$ mass between 3.66 and 3.71 GeV/ c^2 is shown in figure 3(left). An extended unbinned maximum-likelihood fit is performed, where the signal component is modelled by the modified Gaussian function and the background is described by a first order positive polynomial function. The result of this fit is used to obtain the background-subtracted $J/\psi\pi^+\pi^-$ mass distribution from $B^+ \rightarrow J/\psi\pi^+\pi^-K^+$ decays. This distribution is shown in figure 3(right). The yield of the $B^+ \rightarrow (\psi(2S) \rightarrow J/\psi\pi^+\pi^-)K^+$ signal candidates is determined using an unbinned fit to this distribution with a two-component function. The component corresponding to the $B^+ \rightarrow \psi(2S)K^+$ decays is parameterised by the modified Gaussian function. The component describing the $B^+ \rightarrow J/\psi\pi^+\pi^-K^+$ decays without intermediate $\psi(2S)$ state is modelled by a phase-space function,³ modified by a first order positive polynomial function. The tail and resolution parameters for the signal component are taken from simulation with the resolution further corrected by a scale factor, $s_{\psi(2S)}$, that accounts for a small discrepancy between data and simulation [26, 103]. The fit is performed simultaneously to data and simulated samples, as for the signal mode described

³The phase-space mass distribution of a k -body combination of particles from an n -body decay is approximated by $\Phi_{k,n}(x) \propto x_*^{(3k-5)/2} (1-x_*)^{3(n-k)/2-1}$, where $x_* \equiv (x - x_{\min})/(x_{\max} - x_{\min})$, and $x_{\min}$, $x_{\max}$ denote the minimal and maximal values of x , respectively [102]. Here, $k = 3$ and $n = 4$ are used.




Figure 3. Left: mass distributions for selected $B^+ \rightarrow (\psi(2S) \rightarrow J/\psi\pi^+\pi^-)K^+$ decays, with $J/\psi\pi^+\pi^-$ mass between 3.66 and 3.71 GeV/c^2 . Right: background-subtracted $J/\psi\pi^+\pi^-$ mass distribution from selected B^+ decays. The resulting fit functions are overlaid with data distributions.

in section 4. From this fit the number of $B^+ \rightarrow \psi(2S)K^+$ decays is found to be

$$N_{B^+ \rightarrow \psi(2S)K^+} = (121.40 \pm 0.14) \times 10^3, \quad (5.1)$$

where the uncertainty is statistical only. The correction factor $s_{\psi(2S)}$ is found to be 1.057 ± 0.008 , which is in good agreement with results from refs. [15, 26, 98, 99, 103, 104].

6 Branching fraction ratio computation

The ratio of the branching fractions $\mathcal{R}$, defined in eq. (1.1), is calculated as

$$\mathcal{R} = \frac{N_{B^+ \rightarrow J/\psi\eta'K^+}}{N_{B^+ \rightarrow \psi(2S)K^+}} \times \frac{\mathcal{B}(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)}{\mathcal{B}(\eta' \rightarrow \rho^0\gamma)} \times \frac{\epsilon_{B^+ \rightarrow \psi(2S)K^+}}{\epsilon_{B^+ \rightarrow J/\psi\eta'K^+}},$$

where $N_{B^+ \rightarrow J/\psi\eta'K^+}$ and $N_{B^+ \rightarrow \psi(2S)K^+}$ are the yields from eqs. (4.1a) and (5.1), and $\epsilon_{B^+ \rightarrow J/\psi\eta'K^+}$ and $\epsilon_{B^+ \rightarrow \psi(2S)K^+}$ are the efficiencies to reconstruct the observed final states. The efficiencies are the products of detector acceptance, reconstruction, selection and trigger efficiencies, and are calculated using simulated samples, calibrated to match the data as described in section 2. The branching fractions for the $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$ and $\eta' \rightarrow \rho^0\gamma$ decays, $\mathcal{B}(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (34.68 \pm 0.30)\%$ and $\mathcal{B}(\eta' \rightarrow \rho^0\gamma) = (29.4 \pm 0.4)\%$, are taken from ref. [37]. The ratio of branching fractions $\mathcal{R}$ is found to be

$$\frac{\mathcal{B}(B^+ \rightarrow J/\psi\eta'K^+)}{\mathcal{B}(B^+ \rightarrow \psi(2S)K^+)} = (4.91 \pm 0.47) \times 10^{-2},$$

where the uncertainty is statistical only.

As a cross-check, the ratio $\mathcal{R}$ is also calculated for the $\eta' \rightarrow \eta\pi^+\pi^-$, using the signal yields from eqs. (4.1b) and (5.1). The ratio of branching fractions $\mathcal{R}$ is found to be $(4.98 \pm 0.61) \times 10^{-2}$, where the uncertainty is statistical only. This value is in good agreement with that calculated for the $\eta' \rightarrow \rho^0\gamma$ case.

7 Systematic uncertainties

The signal and normalisation channels share the same set of final-state charged particles, and the same trigger and preselection requirements are applied to both. This allows many systematic uncertainties to cancel in the ratio $\mathcal{R}$. The remaining nonnegligible contributions are listed in table 1.

Several systematic uncertainties are associated with the corrections applied to the simulation. The finite size of the $B^+ \rightarrow J/\psi K^+$ signal sample used for correction of the simulated transverse momentum and rapidity spectra of B^+ mesons induces an uncertainty on the B^+ meson p_T and y spectra. In turn, this uncertainty induces small changes in the ratio of efficiencies. The corresponding spread of these changes amounts to 0.1% and is taken as the systematic uncertainty related to the B^+ meson kinematic.

The decay model corrections for the $B^+ \rightarrow J/\psi \eta' K^+$ decay are obtained using the algorithm described in ref. [75]. The systematic uncertainty related to the correction method is estimated by varying the configuration parameters of the algorithm. The largest deviation of the efficiency value from the baseline tuning is found to be 1.1%, which is assigned as systematic uncertainty associated with the B^+ decay model.

There are residual differences in the reconstruction efficiency of charged-particle tracks that do not cancel completely in the ratio of total efficiencies given the slightly different kinematic distributions of the final-state particles. The track-finding efficiencies obtained from simulated samples are corrected using calibration channels [77]. The uncertainties related to the efficiency correction factors are propagated to the ratios of the total efficiencies using pseudoexperiments, and are found to be 0.7%. This value is taken as the systematic uncertainty due to the tracking efficiency calibration.

Differences in the photon reconstruction efficiencies between data and simulation are studied using a large sample of $B^+ \rightarrow J/\psi K^{*+}$ decays, reconstructed using the $K^{*+} \rightarrow K^+ (\pi^0 \rightarrow \gamma\gamma)$ decay mode [78–81]. The uncertainty due to the finite size of the sample is propagated to the ratio of the total efficiencies using pseudoexperiments and is found to be less than 1.0%. The uncertainty due to the accuracy of the $B^+ \rightarrow J/\psi K^{*+}$ branching fraction [37] is 3.5%. These two values are added in quadrature to obtain a systematic uncertainty related to the photon reconstruction of 3.6%.

The kaon identification variable used for the MLP estimator is drawn from calibration data samples and has a dependence on the particle kinematics and track multiplicity. Systematic uncertainties in this procedure arise from the limited size of both the simulation and calibration samples, and the modelling of the particle identification variable. The limitations due to the size of the simulation and calibration samples are evaluated by using bootstrapping techniques [105, 106], creating multiple samples and repeating the procedure for each of these. The impact of potential mismodelling of the kaon identification variable is evaluated by describing the corresponding distributions using density estimates with different kernel widths [76, 107]. For each of these cases, alternative efficiency maps are produced to determine the associated uncertainties. A systematic uncertainty of 2.8% is assigned from the observed differences with alternative efficiency maps.

A systematic uncertainty related to the knowledge of the trigger efficiencies has been previously studied using large samples of $B^+ \rightarrow (J/\psi \rightarrow \mu^+ \mu^-) K^+$ and $B^+ \rightarrow (\psi(2S) \rightarrow \mu^+ \mu^-) K^+$ decays by comparing the ratios of the trigger efficiencies in data and simulation [108]. Based on this comparison, a relative uncertainty of 1.1% is assigned.

The remaining inconsistency between data and simulation, not covered by the corrections discussed in section 2, is estimated by varying the requirement on the response of the MLP classifier in ranges that lead to changes in the measured signal yields as large as $\pm 20\%$. The resulting difference in the data-simulation efficiency ratio is found to be 3.0%.

A different class of systematic uncertainties directly affects the fit itself, namely uncertainties associated with the fit models used to describe the $J/\psi \eta' K^+$, $\psi(2S) K^+$ and $J/\psi \pi^+ \pi^-$ spectra. The systematic uncertainty is accounted for by fits with alternative models. The list of alternative models to describe signal B^+ and $\psi(2S)$ components includes a modified Apollonios function [109], which has exponential instead of power-law tails, a generalised Student's t -distribution [110, 111] and a modified Novosibirsk function [112]. For the combinatorial background for both the signal and the normalisation channels, a third order polynomial function and an exponential function multiplied by a first order polynomial function are chosen as alternative models. For each fit only one component (either signal or background) is replaced at a time, and the same fit function is used for both signal and normalisation channel. The largest deviation of the ratio of signal yields is found to be 0.6%. As alternative background models for $J/\psi \pi^+ \pi^-$ candidates, a first order polynomial function and a product of an exponential function with a first order polynomial function are used. The largest deviation of the signal yield is found to be 1.5%. The two deviations are added in quadrature to obtain a 1.6% systematic uncertainty due to imperfect knowledge of the signal and background shapes.

Finally, the finite size of the simulation samples contributes an uncertainty of 0.9% on the ratio of total efficiencies. The total systematic uncertainty for the ratio of branching fractions $\mathcal{R}$ is calculated as the sum in quadrature of all the values listed above and is found to be 6.0%.

The statistical significance for the $B^+ \rightarrow J/\psi \eta' K^+$ decay is recalculated using Wilks' theorem for each alternative fit model, and the smallest values of 17 and 10 standard deviations for the $\eta' \rightarrow \rho^0 \gamma$ and $\eta' \rightarrow \eta \pi^+ \pi^-$ cases, respectively, are taken as the significance including the systematic uncertainty.

8 Results and summary

The $B^+ \rightarrow J/\psi \eta' K^+$ decays are observed for the first time using proton-proton collision data collected by the LHCb experiment at centre-of-mass energies of 7, 8, and 13 TeV, corresponding to a total integrated luminosity of 9 fb^{-1} . In the analysis, the η' meson is reconstructed from the two $\rho^0 \gamma$ and $\eta \pi^+ \pi^-$ final states. The signal significance exceeds 10 standard deviations for both modes. The branching fraction of the $B^+ \rightarrow J/\psi \eta' K^+$ decay is measured for the $\eta' \rightarrow \rho^0 \gamma$ sample through normalisation to the known branching fraction

Source	Value [%]
B^+ kinematics	0.1
B^+ decay model	1.1
Tracking efficiency correction	0.7
Photon reconstruction correction	3.6
Kaon identification	2.8
Trigger efficiency	1.1
Data-simulation agreement	3.0
Fit model	1.6
Simulation sample size	0.9
Total	6.0

Table 1. Summary of systematic uncertainties on the ratio of branching fractions $\mathcal{R}$. The overall systematic uncertainty is calculated as a sum in quadrature of all the sources.

of the $B^+ \rightarrow \psi(2S)K^+$ decay [37]. The ratio of branching fractions is found to be

$$\frac{\mathcal{B}(B^+ \rightarrow J/\psi\eta'K^+)}{\mathcal{B}(B^+ \rightarrow \psi(2S)K^+)} = (4.91 \pm 0.47 \pm 0.29 \pm 0.07) \times 10^{-2},$$

where the first uncertainty is statistical, the second is systematic and the third is related to the uncertainties on the branching fractions of the intermediate resonances. The absolute branching fraction is determined using the known branching fraction of $B^+ \rightarrow \psi(2S)K^+$ decays, $\mathcal{B}(B^+ \rightarrow \psi(2S)K^+) = (6.24 \pm 0.20) \times 10^{-4}$ [37], and is found to be

$$\mathcal{B}(B^+ \rightarrow J/\psi\eta'K^+) = (3.06 \pm 0.29 \pm 0.18 \pm 0.04) \times 10^{-5},$$

where the first uncertainty is statistical, the second is systematic and the third is due to external branching fractions uncertainties. The measured branching fraction is consistent with the upper limit previously set by the Belle collaboration [58]. An inspection of the $J/\psi\eta'$ mass spectrum shows no significant contributions from the decays via intermediate charmonium or charmonium-like resonances.

Acknowledgments

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MICINN (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), Polish WLCG (Poland) and

NERSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from ARC and ARDC (Australia); Minciencias (Colombia); AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); A*MIDEX, ANR, IPhU and Labex P2IO, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, CAS CCEPP, Fundamental Research Funds for the Central Universities, and Sci. & Tech. Program of Guangzhou (China); GVA, XuntaGal, GENCAT and Prog. Atracción Talento, CM (Spain); SRC (Sweden); the Leverhulme Trust, the Royal Society and UKRI (United Kingdom).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- [1] BELLE collaboration, *Observation of a narrow charmonium-like state in exclusive $B^\pm \rightarrow K^\pm \pi^+ \pi^- J/\psi$ decays*, *Phys. Rev. Lett.* **91** (2003) 262001 [[hep-ex/0309032](#)] [[INSPIRE](#)].
- [2] LHCb collaboration, *Observation of $J/\psi p$ resonances consistent with pentaquark states in $\Lambda_b^0 \rightarrow J/\psi K^- p$ decays*, *Phys. Rev. Lett.* **115** (2015) 072001 [LHCb-PAPER-2015-029, CERN-PH-EP-2015-153] [[arXiv:1507.03414](#)] [[INSPIRE](#)].
- [3] LHCb collaboration, *Model-independent evidence for $J/\psi p$ contributions to $\Lambda_b^0 \rightarrow J/\psi p K^-$ decays*, *Phys. Rev. Lett.* **117** (2016) 082002 [LHCb-PAPER-2016-009, CERN-EP-2016-086] [[arXiv:1604.05708](#)] [[INSPIRE](#)].
- [4] LHCb collaboration, *Observation of a narrow pentaquark state, $P_c(4312)^+$, and of two-peak structure of the $P_c(4450)^+$* , *Phys. Rev. Lett.* **122** (2019) 222001 [LHCb-PAPER-2019-014, CERN-EP-2019-058] [[arXiv:1904.03947](#)] [[INSPIRE](#)].
- [5] LHCb collaboration, *Evidence for exotic hadron contributions to $\Lambda_b^0 \rightarrow J/\psi p \pi^-$ decays*, *Phys. Rev. Lett.* **117** (2016) 082003 [Addendum *ibid.* **117** (2016) 109902] [LHCb-PAPER-2016-015, CERN-EP-2016-151] [[arXiv:1606.06999](#)] [[INSPIRE](#)].
- [6] LHCb collaboration, *Evidence for a new structure in the $J/\psi p$ and $J/\psi \bar{p}$ systems in $B_s^0 \rightarrow J/\psi p \bar{p}$ decays*, *Phys. Rev. Lett.* **128** (2022) 062001 [LHCb-PAPER-2021-018, CERN-EP-2021-150] [[arXiv:2108.04720](#)] [[INSPIRE](#)].
- [7] LHCb collaboration, *Evidence of a $J/\psi \Lambda$ structure and observation of excited Ξ^- states in the $\Xi_b^- \rightarrow J/\psi \Lambda K^-$ decay*, *Sci. Bull.* **66** (2021) 1278 [LHCb-PAPER-2020-039, CERN-EP-2020-233] [[arXiv:2012.10380](#)] [[INSPIRE](#)].
- [8] BELLE collaboration, *Observation of a resonance-like structure in the $\pi^\pm \psi'$ mass distribution in exclusive $B \rightarrow K \pi^\pm \psi'$ decays*, *Phys. Rev. Lett.* **100** (2008) 142001 [[arXiv:0708.1790](#)] [[INSPIRE](#)].

- [9] BELLE collaboration, *Dalitz analysis of $B \rightarrow K\pi^+\psi'$ decays and the $Z(4430)^+$* , *Phys. Rev. D* **80** (2009) 031104 [[arXiv:0905.2869](#)] [[INSPIRE](#)].
- [10] BELLE collaboration, *Experimental constraints on the spin and parity of the $Z(4430)^+$* , *Phys. Rev. D* **88** (2013) 074026 [[arXiv:1306.4894](#)] [[INSPIRE](#)].
- [11] LHCb collaboration, *Observation of the resonant character of the $Z(4430)^-$ state*, *Phys. Rev. Lett.* **112** (2014) 222002 [LHCb-PAPER-2014-014, CERN-PH-EP-2014-061] [[arXiv:1404.1903](#)] [[INSPIRE](#)].
- [12] LHCb collaboration, *Model-independent confirmation of the $Z(4430)^-$ state*, *Phys. Rev. D* **92** (2015) 112009 [LHCb-PAPER-2015-038, CERN-PH-EP-2015-244] [[arXiv:1510.01951](#)] [[INSPIRE](#)].
- [13] LHCb collaboration, *Observation of $J/\psi\phi$ structures consistent with exotic states from amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ decays*, *Phys. Rev. Lett.* **118** (2017) 022003 [LHCb-PAPER-2016-018, CERN-EP-2016-155] [[arXiv:1606.07895](#)] [[INSPIRE](#)].
- [14] LHCb collaboration, *Amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ decays*, *Phys. Rev. D* **95** (2017) 012002 [LHCb-PAPER-2016-019, CERN-EP-2016-156] [[arXiv:1606.07898](#)] [[INSPIRE](#)].
- [15] LHCb collaboration, *Study of $B_s^0 \rightarrow J\psi\pi^+\pi^-K^+K^-$ decays*, *JHEP* **02** (2021) 024 [LHCb-PAPER-2020-035, CERN-EP-2020-192] [*Erratum ibid.* **04** (2021) 170] [[arXiv:2011.01867](#)] [[INSPIRE](#)].
- [16] LHCb collaboration, *Observation of new resonances decaying to $J/\psi K^+$ and $J/\psi\phi$* , *Phys. Rev. Lett.* **127** (2021) 082001 [LHCb-PAPER-2020-044, CERN-EP-2021-025] [[arXiv:2103.01803](#)] [[INSPIRE](#)].
- [17] LHCb collaboration, *Evidence of a $J/\psi K_S^0$ structure in $B^0 \rightarrow J/\psi\phi K_S^0$ decays*, [arXiv:2301.04899](#) [LHCb-PAPER-2022-040, CERN-EP-2022-258] [[INSPIRE](#)].
- [18] LHCb collaboration, *Evidence for an $\eta_c(1S)\pi^-$ resonance in $B^0 \rightarrow \eta_c(1S)K^+\pi^-$ decays*, *Eur. Phys. J. C* **78** (2018) 1019 [LHCb-PAPER-2018-034, CERN-EP-2018-245] [[arXiv:1809.07416](#)] [[INSPIRE](#)].
- [19] LHCb collaboration, *Model-independent observation of exotic contributions to $B^0 \rightarrow J/\psi K^+\pi^-$ decays*, *Phys. Rev. Lett.* **122** (2019) 152002 [LHCb-PAPER-2018-043, CERN-EP-2018-330] [[arXiv:1901.05745](#)] [[INSPIRE](#)].
- [20] LHCb collaboration, *Observation of $B_s^0 \rightarrow \chi_{c1}\phi$ decay and study of $B^0 \rightarrow \chi_{c1,2}K^{*0}$ decays*, *Nucl. Phys. B* **874** (2013) 663 [LHCb-PAPER-2013-024, CERN-PH-EP-2013-088] [[arXiv:1305.6511](#)] [[INSPIRE](#)].
- [21] LHCb collaboration, *Evidence for the decay $X(3872) \rightarrow \psi(2S)\gamma$* , *Nucl. Phys. B* **886** (2014) 665 [LHCb-PAPER-2014-008, CERN-PH-EP-2014-050] [[arXiv:1404.0275](#)] [[INSPIRE](#)].
- [22] LHCb collaboration, *Observation of the decays $\Lambda_b^0 \rightarrow \chi_{c1}pK^-$ and $\Lambda_b^0 \rightarrow \chi_{c2}pK^-$* , *Phys. Rev. Lett.* **119** (2017) 062001 [LHCb-PAPER-2017-011, CERN-EP-2017-073] [[arXiv:1704.07900](#)] [[INSPIRE](#)].
- [23] LHCb collaboration, *Observation of the decay $\Lambda_b^0 \rightarrow \chi_{c1}p\pi^-$* , *JHEP* **05** (2021) 095 [LHCb-PAPER-2021-003, CERN-EP-2021-037] [[arXiv:2103.04949](#)] [[INSPIRE](#)].

- [24] LHCb collaboration, *Determination of the $X(3872)$ meson quantum numbers*, *Phys. Rev. Lett.* **110** (2013) 222001 [LHCb-PAPER-2013-001, CERN-PH-EP-2013-017] [[arXiv:1302.6269](#)] [[INSPIRE](#)].
- [25] LHCb collaboration, *Quantum numbers of the $X(3872)$ state and orbital angular momentum in its $\rho^0 J/\psi$ decay*, *Phys. Rev. D* **92** (2015) 011102 [LHCb-PAPER-2015-015, CERN-PH-EP-2015-098] [[arXiv:1504.06339](#)] [[INSPIRE](#)].
- [26] LHCb collaboration, *Study of the $\psi_2(3823)$ and $\chi_{c1}(3872)$ states in $B^+ \rightarrow (J/\psi \pi^+ \pi^-) K^+$ decays*, *JHEP* **08** (2020) 123 [LHCb-PAPER-2020-009, CERN-EP-2020-071] [[arXiv:2005.13422](#)] [[INSPIRE](#)].
- [27] LHCb collaboration, *Observation of sizeable ω contribution to $\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi$ decays*, *Phys. Rev. D* **108** (2023) L011103 [LHCb-PAPER-2021-045] [[arXiv:2204.12597](#)] [[INSPIRE](#)].
- [28] BABAR collaboration, *Evidence for the decay $X(3872) \rightarrow J/\psi \omega$* , *Phys. Rev. D* **82** (2010) 011101 [[arXiv:1005.5190](#)] [[INSPIRE](#)].
- [29] LHCb collaboration, *Study of charmonium and charmonium-like contributions in $B^+ \rightarrow J/\psi \eta K^+$ decays*, *JHEP* **04** (2022) 046 [LHCb-PAPER-2021-047, CERN-EP-2022-009] [[arXiv:2202.04045](#)] [[INSPIRE](#)].
- [30] BELLE collaboration, *Observation of $\psi(4040)$ and $\psi(4160)$ decay into $\eta J/\psi$* , *Phys. Rev. D* **87** (2013) 051101 [[arXiv:1210.7550](#)] [[INSPIRE](#)].
- [31] BESIII collaboration, *Observation of the $Y(4220)$ and $Y(4360)$ in the process $e^+ e^- \rightarrow \eta J/\psi$* , *Phys. Rev. D* **102** (2020) 031101 [[arXiv:2003.03705](#)] [[INSPIRE](#)].
- [32] N. Brambilla et al., *The XYZ states: experimental and theoretical status and perspectives*, *Phys. Rept.* **873** (2020) 1 [[arXiv:1907.07583](#)] [[INSPIRE](#)].
- [33] A. Ali, L. Maiani and A.D. Polosa, *Multiquark hadrons*, Cambridge University Press (2019) [[DOI:10.1017/9781316761465](#)] [[INSPIRE](#)].
- [34] L. Maiani and A. Pilloni, *GGI Lectures on exotic hadrons*, [arXiv:2207.05141](#) [[INSPIRE](#)].
- [35] N. Brambilla et al., *Substructure of multiquark hadrons (Snowmass 2021 White Paper)*, [arXiv:2203.16583](#) [[INSPIRE](#)].
- [36] N. Brambilla et al., *Heavy Quarkonium: progress, puzzles, and opportunities*, *Eur. Phys. J. C* **71** (2011) 1534 [[arXiv:1010.5827](#)] [[INSPIRE](#)].
- [37] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *PTEP* **2022** (2022) 083C01 [[INSPIRE](#)].
- [38] J.L. Rosner, *Quark content of neutral mesons*, *Phys. Rev. D* **27** (1983) 1101 [[INSPIRE](#)].
- [39] A. Bramon, R. Escribano and M.D. Scadron, *The $\eta - \eta'$ mixing angle revisited*, *Eur. Phys. J. C* **7** (1999) 271 [[hep-ph/9711229](#)] [[INSPIRE](#)].
- [40] A. Bramon, R. Escribano and M.D. Scadron, *Mixing of $\eta - \eta'$ mesons in J/ψ decays into a vector and a pseudoscalar meson*, *Phys. Lett. B* **403** (1997) 339 [[hep-ph/9703313](#)] [[INSPIRE](#)].

- [41] V.A. Novikov, M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *A theory of the $J/\psi \rightarrow \eta(\eta')\gamma$ decays*, *Nucl. Phys. B* **165** (1980) 55 [INSPIRE].
- [42] V.A. Novikov, M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *η' meson as pseudoscalar gluonium*, *Phys. Lett. B* **86** (1979) 347 [INSPIRE].
- [43] V.A. Novikov, M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *In a search for scalar gluonium*, *Nucl. Phys. B* **165** (1980) 67 [INSPIRE].
- [44] A.L. Kataev, N.V. Krasnikov and A.A. Pivovarov, *The connection between the scales of the gluon and quark worlds in perturbative QCD*, *Phys. Lett. B* **107** (1981) 115 [INSPIRE].
- [45] A.L. Kataev, N.V. Krasnikov and A.A. Pivovarov, *Two loop calculations for the propagators of gluonic currents*, *Nucl. Phys. B* **198** (1982) 508 [hep-ph/9612326] [INSPIRE].
- [46] C.E. Thomas, *Composition of the pseudoscalar η and η' mesons*, *JHEP* **10** (2007) 026 [arXiv:0705.1500] [INSPIRE].
- [47] R. Escribano and J. Nadal, *On the gluon content of the η and η' mesons*, *JHEP* **05** (2007) 006 [hep-ph/0703187] [INSPIRE].
- [48] R. Escribano, *$J/\psi \rightarrow VP$ decays and the quark and gluon content of the eta and eta-prime*, *Eur. Phys. J. C* **65** (2010) 467 [arXiv:0807.4201] [INSPIRE].
- [49] F. Ambrosino et al., *A Global fit to determine the pseudoscalar mixing angle and the gluonium content of the eta-prime meson*, *JHEP* **07** (2009) 105 [arXiv:0906.3819] [INSPIRE].
- [50] CLEO collaboration, *Absolute branching fraction measurements for exclusive D_s semileptonic decays*, *Phys. Rev. D* **80** (2009) 052007 [arXiv:0903.0601] [INSPIRE].
- [51] CLEO collaboration, *Studies of $D^+ \rightarrow \eta', \eta, \phi e^+ \nu_e$* , *Phys. Rev. D* **84** (2011) 032001 [arXiv:1011.1195] [INSPIRE].
- [52] C. Di Donato, G. Ricciardi and I. Bigi, *$\eta - \eta'$ mixing — from electromagnetic transitions to weak decays of charm and beauty hadrons*, *Phys. Rev. D* **85** (2012) 013016 [arXiv:1105.3557] [INSPIRE].
- [53] LHCb collaboration, *Study of $\eta - \eta'$ mixing from measurement of $B_{(s)}^0 \rightarrow J/\psi \eta^{(\prime)}$ decay rates*, *JHEP* **01** (2015) 024 [LHCb-PAPER-2014-056, CERN-PH-EP-2014-266] [arXiv:1411.0943] [INSPIRE].
- [54] M.A. Andreichikov, M.I. Vysotsky and V.A. Novikov, *On the branching ratios of the $B^0 \rightarrow J/\psi \eta(\eta', \pi^0)$ and $B_s \rightarrow J/\psi \eta(\eta')$ decays*, *Pisma Zh. Eksp. Teor. Fiz.* **110** (2019) 633 [INSPIRE].
- [55] M.A. Andreichikov, M.I. Eides, V.A. Novikov and M.I. Vysotsky, *The physics of the $\eta - \eta'$ system versus $B^0 \rightarrow J/\psi \eta(\eta')$ and $B_s \rightarrow J/\psi \eta(\eta')$ decays*, *Int. J. Mod. Phys. A* **35** (2020) 2050111 [arXiv:1911.10596] [INSPIRE].
- [56] F.E. Close et al., *Gluonic hadrons and charmless B decays*, *Phys. Rev. D* **57** (1998) 5653 [hep-ph/9708265] [INSPIRE].

- [57] F.E. Close and P.R. Page, *Gluonic charmonium resonances at BaBar and BELLE?*, *Phys. Lett. B* **628** (2005) 215 [[hep-ph/0507199](#)] [[INSPIRE](#)].
- [58] BELLE collaboration, *Search for $B^+ \rightarrow J/\psi \eta' K^+$ and $B^0 \rightarrow J/\psi \eta' K_S^0$ decays*, *Phys. Rev. D* **75** (2007) 017101 [[hep-ex/0610084](#)] [[INSPIRE](#)].
- [59] LHCb collaboration, *The LHCb detector at the LHC*, 2008 *JINST* **3** S08005 [[INSPIRE](#)].
- [60] LHCb collaboration, *LHCb Detector Performance*, *Int. J. Mod. Phys. A* **30** (2015) 1530022 [[LHCb-DP-2014-002](#), [CERN-PH-EP-2014-290](#)] [[arXiv:1412.6352](#)] [[INSPIRE](#)].
- [61] R. Aaij et al., *Performance of the LHCb Vertex Locator*, 2014 *JINST* **9** P09007 [[LHCb-DP-2014-001](#)] [[arXiv:1405.7808](#)] [[INSPIRE](#)].
- [62] R. Arink et al., *Performance of the LHCb Outer Tracker*, 2014 *JINST* **9** P01002 [[arXiv:1311.3893](#)] [[INSPIRE](#)].
- [63] P. d'Argent et al., *Improved performance of the LHCb Outer Tracker in LHC Run 2*, 2017 *JINST* **12** P11016 [[arXiv:1708.00819](#)] [[INSPIRE](#)].
- [64] M. Adinolfi et al., *Performance of the LHCb RICH detector at the LHC*, *Eur. Phys. J. C* **73** (2013) 2431 [[arXiv:1211.6759](#)] [[INSPIRE](#)].
- [65] A.A. Alves Jr. et al., *Performance of the LHCb muon system*, 2013 *JINST* **8** P02022 [[arXiv:1211.1346](#)] [[INSPIRE](#)].
- [66] R. Aaij et al., *The LHCb trigger and its performance in 2011*, 2013 *JINST* **8** P04022 [[arXiv:1211.3055](#)] [[INSPIRE](#)].
- [67] LHCb collaboration, *Design and performance of the LHCb trigger and full real-time reconstruction in Run 2 of the LHC*, 2019 *JINST* **14** P04013 [[arXiv:1812.10790](#)] [[INSPIRE](#)].
- [68] T. Sjostrand, S. Mrenna and P.Z. Skands, *A brief introduction to PYTHIA 8.1*, *Comput. Phys. Commun.* **178** (2008) 852 [[arXiv:0710.3820](#)] [[INSPIRE](#)].
- [69] LHCb collaboration, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, *J. Phys. Conf. Ser.* **331** (2011) 032047 [[INSPIRE](#)].
- [70] D.J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152 [[INSPIRE](#)].
- [71] N. Davidson, T. Przydzinski and Z. Was, *PHOTOS interface in C++: technical and physics documentation*, *Comput. Phys. Commun.* **199** (2016) 86 [[arXiv:1011.0937](#)] [[INSPIRE](#)].
- [72] J. Allison et al., *Geant4 developments and applications*, *IEEE Trans. Nucl. Sci.* **53** (2006) 270 [[INSPIRE](#)].
- [73] GEANT4 collaboration, *GEANT4: a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [74] LHCb collaboration, *The LHCb simulation application, Gauss: Design, evolution and experience*, *J. Phys. Conf. Ser.* **331** (2011) 032023 [[INSPIRE](#)].

- [75] A. Rogozhnikov, *Reweighting with Boosted Decision Trees*, *J. Phys. Conf. Ser.* **762** (2016) 012036 [[arXiv:1608.05806](#)] [[INSPIRE](#)].
- [76] R. Aaij et al., *Selection and processing of calibration samples to measure the particle identification performance of the LHCb experiment in Run 2*, *EPJ Tech. Instrum.* **6** (2019) 1 [[arXiv:1803.00824](#)] [[INSPIRE](#)].
- [77] LHCb collaboration, *Measurement of the track reconstruction efficiency at LHCb*, 2015 *JINST* **10** P02007 [CERN-LHCB-DP-2013-002] [[arXiv:1408.1251](#)] [[INSPIRE](#)].
- [78] LHCb collaboration, *Evidence for the decay $B^0 \rightarrow J/\psi\omega$ and measurement of the relative branching fractions of B_s^0 meson decays to $J/\psi\eta$ and $J/\psi\eta'$* , *Nucl. Phys. B* **867** (2013) 547 [LHCb-PAPER-2012-022, CERN-PH-EP-2012-287] [[arXiv:1210.2631](#)] [[INSPIRE](#)].
- [79] LHCb collaboration, *Observations of $B_s^0 \rightarrow \psi(2S)\eta$ and $B_{(s)}^0 \rightarrow \psi(2S)\pi^+\pi^-$ decays*, *Nucl. Phys. B* **871** (2013) 403 [LHCb-PAPER-2012-053, CERN-PH-EP-2013-024] [[arXiv:1302.6354](#)] [[INSPIRE](#)].
- [80] E. Govorkova, *Study of π^0/γ efficiency using B meson decays in the LHCb experiment*, *Phys. Atom. Nucl.* **79** (2016) 1474 [[arXiv:1505.02960](#)] [[INSPIRE](#)].
- [81] K. Govorkova, *Study of photons and neutral pions reconstruction efficiency in the LHCb experiment*, CERN-THESIS-2015-272.
- [82] A. Powell et al., *Particle identification at LHCb*, *PoS ICHEP2010* (2010) 020 [LHCb-PROC-2011-008].
- [83] H. Terrier and I. Belyaev, *Particle identification with LHCb calorimeters*, LHCb-2003-092.
- [84] C. Abellán Beteta et al., *Calibration and performance of the LHCb calorimeters in Run 1 and 2 at the LHC*, [arXiv:2008.11556](#) [[INSPIRE](#)].
- [85] TASSO collaboration, *Measurement of the radiative width of the $\eta'(958)$ in two photon interactions*, *Phys. Lett. B* **147** (1984) 487 [[INSPIRE](#)].
- [86] H. Kolanoski, *Two-photon physics at e^+e^- storage rings*, vol. 105, Berlin, Springer-Verlag (1984) [[DOI:10.1007/BFb0045900](#)] [[INSPIRE](#)].
- [87] ARGUS collaboration, *Observation of the Decay $D_s^+ \rightarrow \eta'\pi^+$* , *Phys. Lett. B* **245** (1990) 315 [[INSPIRE](#)].
- [88] CLEO collaboration, *Two-body D_s^+ decays to $\eta\pi^+$, $\eta'\pi^+$, $\eta\rho^+$, $\eta'\rho^+$ and $\phi\rho^+$* , *Phys. Rev. D* **45** (1992) 3965 [[INSPIRE](#)].
- [89] CLEO collaboration, *D_s^+ decays to $\eta\pi^+$ and $\eta'\pi^+$* , *Phys. Rev. Lett.* **68** (1992) 1275 [[INSPIRE](#)].
- [90] BESIII collaboration, *Precision study of $\eta' \rightarrow \gamma\pi^+\pi^-$ decay dynamics*, *Phys. Rev. Lett.* **120** (2018) 242003 [[arXiv:1712.01525](#)] [[INSPIRE](#)].
- [91] W.D. Hulsbergen, *Decay chain fitting with a Kalman filter*, *Nucl. Instrum. Meth. A* **552** (2005) 566 [[physics/0503191](#)] [[INSPIRE](#)].

- [92] W.S. McCulloch and W. Pitts, *A logical calculus of the ideas immanent in nervous activity*, *The Bulletin of Mathematical Biophysics* **5** (1943) 115.
- [93] F. Rosenblatt, *The perceptron: a probabilistic model for information storage and organization in the brain*, *Psychological Review* **65** (1958) 386.
- [94] J.-H. Zhong et al., *A program for the Bayesian neural network in the ROOT framework*, *Comput. Phys. Commun.* **182** (2011) 2655 [[arXiv:1103.2854](#)] [[INSPIRE](#)].
- [95] T. Skwarnicki, *A study of the radiative CASCADE transitions between the Upsilon-Prime and Upsilon resonances*, Ph.D. thesis, Institute of Nuclear Physics, Krakow (1986) [[INSPIRE](#)].
- [96] LHCb collaboration, *Observation of J/ψ pair production in pp collisions at $\sqrt{s} = 7\text{TeV}$* , *Phys. Lett. B* **707** (2012) 52 [LHCb-PAPER-2011-013, CERN-PH-EP-2011-135] [[arXiv:1109.0963](#)] [[INSPIRE](#)].
- [97] S. Karlin and L.S. Shapley, *Geometry of moment spaces*, vol. 12 of *Memoirs of the American Mathematical Society*, American Mathematical Society, Providence, Rhode Island (1953).
- [98] LHCb collaboration, *Study of B_c^+ decays to charmonia and three light hadrons*, *JHEP* **01** (2022) 065 [LHCb-PAPER-2021-034, CERN-EP-2021-216] [[arXiv:2111.03001](#)] [[INSPIRE](#)].
- [99] LHCb collaboration, *Study of B_c^+ meson decays to charmonia plus multihadron final states*, [arXiv:2208.08660](#) [LHCb-PAPER-2022-025, CERN-EP-2022-162] [[INSPIRE](#)].
- [100] S.S. Wilks, *The large-sample distribution of the likelihood ratio for testing composite hypotheses*, *Annals Math. Statist.* **9** (1938) 60 [[INSPIRE](#)].
- [101] M. Pivk and F.R. Le Diberder, *SPlot: a statistical tool to unfold data distributions*, *Nucl. Instrum. Meth. A* **555** (2005) 356 [[physics/0402083](#)] [[INSPIRE](#)].
- [102] E. Byckling and K. Kajantie, *Particle kinematics*, John Wiley & Sons Inc., New York (1973).
- [103] LHCb collaboration, *Study of the lineshape of the $\chi_{c1}(3872)$ state*, *Phys. Rev. D* **102** (2020) 092005 [LHCb-PAPER-2020-008, CERN-EP-2020-086] [[arXiv:2005.13419](#)] [[INSPIRE](#)].
- [104] LHCb collaboration, *Observation of the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ decay*, [arXiv:2302.10629](#) [[INSPIRE](#)].
- [105] B. Efron, *Bootstrap methods: another look at the Jackknife*, *Annals Statist.* **7** (1979) 1 [[INSPIRE](#)].
- [106] B. Efron and R.J. Tibshirani, *An introduction to the bootstrap*, Chapman and Hall, New York (1994) [[DOI:10.1201/9780429246593](#)].
- [107] A. Poluektov, *Kernel density estimation of a multidimensional efficiency profile*, 2015 *JINST* **10** P02011 [[arXiv:1411.5528](#)] [[INSPIRE](#)].
- [108] LHCb collaboration, *Measurement of relative branching fractions of B decays to $\psi(2S)$ and J/ψ mesons*, *Eur. Phys. J. C* **72** (2012) 2118 [LHCb-PAPER-2012-010, CERN-PH-EP-2012-113] [[arXiv:1205.0918](#)] [[INSPIRE](#)].

- [109] D. Martínez Santos and F. Dupertuis, *Mass distributions marginalized over per-event errors*, *Nucl. Instrum. Meth. A* **764** (2014) 150 [[arXiv:1312.5000](#)] [[INSPIRE](#)].
- [110] Student, *The probable error of a mean*, *Biometrika* **6** (1908) 1.
- [111] S. Jackman, *Bayesian analysis for the social sciences*, John Wiley & Sons, Inc., Hoboken, New Jersey, U.S.A. (2009).
- [112] BABAR collaboration, *Branching fraction measurements of the color-suppressed decays $\bar{B}^0 \rightarrow D^{(*)0}\pi^0$, $D^{(*)0}\eta$, $D^{(*)0}\omega$, and $D^{(*)0}\eta'$ and measurement of the polarization in the decay $\bar{B}^0 \rightarrow D^{*0}\omega$* , *Phys. Rev. D* **84** (2011) 112007 [Erratum *ibid.* **87** (2013) 039901] [[arXiv:1107.5751](#)] [[INSPIRE](#)].

The LHCb collaboration

R. Aaij ^{[ID](#)32}, A.S.W. Abdelmotteleb ^{[ID](#)51}, C. Abellan Beteta ^{[ID](#)45}, F. Abudinén ^{[ID](#)51}, T. Ackernley ^{[ID](#)55}, B. Adeva ^{[ID](#)41}, M. Adinolfi ^{[ID](#)49}, P. Adlarson ^{[ID](#)77}, H. Afsharnia ^{[ID](#)9}, C. Agapopoulou ^{[ID](#)13}, C.A. Aidala ^{[ID](#)78}, Z. Ajaltouni ^{[ID](#)9}, S. Akar ^{[ID](#)60}, K. Akiba ^{[ID](#)32}, P. Albicocco ^{[ID](#)23}, J. Albrecht ^{[ID](#)15}, F. Alessio ^{[ID](#)43}, M. Alexander ^{[ID](#)54}, A. Alfonso Alberio ^{[ID](#)40}, Z. Aliouche ^{[ID](#)57}, P. Alvarez Cartelle ^{[ID](#)50}, R. Amalric ^{[ID](#)13}, S. Amato ^{[ID](#)2}, J.L. Amey ^{[ID](#)49}, Y. Amhis ^{[ID](#)11,43}, L. An ^{[ID](#)43}, L. Anderlini ^{[ID](#)22}, M. Andersson ^{[ID](#)45}, A. Andreianov ^{[ID](#)38}, M. Andreotti ^{[ID](#)21}, D. Andreou ^{[ID](#)63}, D. Ao ^{[ID](#)6}, F. Archilli ^{[ID](#)31,t}, A. Artamonov ^{[ID](#)38}, M. Artuso ^{[ID](#)63}, E. Aslanides ^{[ID](#)10}, M. Atzeni ^{[ID](#)45}, B. Audurier ^{[ID](#)12}, I. Bachiller Perea ^{[ID](#)8}, S. Bachmann ^{[ID](#)17}, M. Bachmayer ^{[ID](#)44}, J.J. Back ^{[ID](#)51}, A. Bailly-reyre ^{[ID](#)13}, P. Baladron Rodriguez ^{[ID](#)41}, V. Balagura ^{[ID](#)12}, W. Baldini ^{[ID](#)21,43}, J. Baptista de Souza Leite ^{[ID](#)1}, M. Barbeti ^{[ID](#)22,k}, R.J. Barlow ^{[ID](#)57}, S. Barsuk ^{[ID](#)11}, W. Barter ^{[ID](#)53}, M. Bartolini ^{[ID](#)50}, F. Baryshnikov ^{[ID](#)38}, J.M. Basels ^{[ID](#)14}, G. Bassi ^{[ID](#)29,q}, B. Batsukh ^{[ID](#)4}, A. Battig ^{[ID](#)15}, A. Bay ^{[ID](#)44}, A. Beck ^{[ID](#)51}, M. Becker ^{[ID](#)15}, F. Bedeschi ^{[ID](#)29}, I.B. Bediaga ^{[ID](#)1}, A. Beiter ^{[ID](#)63}, S. Belin ^{[ID](#)41}, V. Bellee ^{[ID](#)45}, K. Belous ^{[ID](#)38}, I. Belov ^{[ID](#)38}, I. Belyaev ^{[ID](#)38}, G. Benane ^{[ID](#)10}, G. Bencivenni ^{[ID](#)23}, E. Ben-Haim ^{[ID](#)13}, A. Berezhniov ^{[ID](#)38}, R. Bernet ^{[ID](#)45}, S. Bernet Andres ^{[ID](#)39}, D. Berninghoff ^{[ID](#)17}, H.C. Bernstein ^{[ID](#)63}, C. Bertella ^{[ID](#)57}, A. Bertolin ^{[ID](#)28}, C. Betancourt ^{[ID](#)45}, F. Betti ^{[ID](#)43}, Ia. Bezshyiko ^{[ID](#)45}, J. Bhom ^{[ID](#)35}, L. Bian ^{[ID](#)69}, M.S. Bieker ^{[ID](#)15}, N.V. Biesuz ^{[ID](#)21}, P. Billoir ^{[ID](#)13}, A. Biolchini ^{[ID](#)32}, M. Birch ^{[ID](#)56}, F.C.R. Bishop ^{[ID](#)50}, A. Bitadze ^{[ID](#)57}, A. Bizzeti ^{[ID](#)}, M.P. Blago ^{[ID](#)50}, T. Blake ^{[ID](#)51}, F. Blanc ^{[ID](#)44}, J.E. Blank ^{[ID](#)15}, S. Blusk ^{[ID](#)63}, D. Bobulska ^{[ID](#)54}, V. Bocharnikov ^{[ID](#)38}, J.A. Boelhauve ^{[ID](#)15}, O. Boente Garcia ^{[ID](#)12}, T. Boettcher ^{[ID](#)60}, A. Boldyrev ^{[ID](#)38}, C.S. Bolognani ^{[ID](#)75}, R. Bolzonella ^{[ID](#)21,j}, N. Bondar ^{[ID](#)38,43}, F. Borgato ^{[ID](#)28}, S. Borghi ^{[ID](#)57}, M. Borsato ^{[ID](#)17}, J.T. Borsuk ^{[ID](#)35}, S.A. Bouchiba ^{[ID](#)44}, T.J.V. Bowcock ^{[ID](#)55}, A. Boyer ^{[ID](#)43}, C. Bozzi ^{[ID](#)21}, M.J. Bradley ^{[ID](#)56}, S. Braun ^{[ID](#)61}, A. Brea Rodriguez ^{[ID](#)41}, N. Breer ^{[ID](#)15}, J. Brodzicka ^{[ID](#)35}, A. Brossa Gonzalo ^{[ID](#)41}, J. Brown ^{[ID](#)55}, D. Brundu ^{[ID](#)27}, A. Buonauro ^{[ID](#)45}, L. Buonincontri ^{[ID](#)28}, A.T. Burke ^{[ID](#)57}, C. Burr ^{[ID](#)43}, A. Bursche ^{[ID](#)67}, A. Butkevich ^{[ID](#)38}, J.S. Butter ^{[ID](#)32}, J. Buytaert ^{[ID](#)43}, W. Byczynski ^{[ID](#)43}, S. Cadeddu ^{[ID](#)27}, H. Cai ^{[ID](#)69}, R. Calabrese ^{[ID](#)21,j}, L. Calefice ^{[ID](#)15}, S. Cali ^{[ID](#)23}, M. Calvi ^{[ID](#)26,n}, M. Calvo Gomez ^{[ID](#)39}, P. Campana ^{[ID](#)23}, D.H. Campora Perez ^{[ID](#)75}, A.F. Campoverde Quezada ^{[ID](#)6}, S. Capelli ^{[ID](#)26,n}, L. Capriotti ^{[ID](#)20}, A. Carbone ^{[ID](#)20,h}, R. Cardinale ^{[ID](#)24,l}, A. Cardini ^{[ID](#)27}, P. Carniti ^{[ID](#)26,n}, L. Carus ^{[ID](#)14}, A. Casais Vidal ^{[ID](#)41}, R. Caspary ^{[ID](#)17}, G. Casse ^{[ID](#)55}, M. Cattaneo ^{[ID](#)43}, G. Cavallero ^{[ID](#)21}, V. Cavallini ^{[ID](#)21,j}, S. Celani ^{[ID](#)44}, J. Cerasoli ^{[ID](#)10}, D. Cervenkov ^{[ID](#)58}, A.J. Chadwick ^{[ID](#)55}, I. Chahrour ^{[ID](#)78}, M.G. Chapman ^{[ID](#)49}, M. Charles ^{[ID](#)13}, Ph. Charpentier ^{[ID](#)43}, C.A. Chavez Barajas ^{[ID](#)55}, M. Chefdeville ^{[ID](#)8}, C. Chen ^{[ID](#)10}, S. Chen ^{[ID](#)4}, A. Chernov ^{[ID](#)35}, S. Chernyshenko ^{[ID](#)47}, V. Chobanova ^{[ID](#)41}, S. Cholak ^{[ID](#)44}, M. Chrzasczcz ^{[ID](#)35}, A. Chubykin ^{[ID](#)38}, V. Chulikov ^{[ID](#)38}, P. Ciambrone ^{[ID](#)23}, M.F. Cicala ^{[ID](#)51}, X. Cid Vidal ^{[ID](#)41}, G. Ciezarek ^{[ID](#)43}, P. Cifra ^{[ID](#)43}, P.E.L. Clarke ^{[ID](#)53}, M. Clemencic ^{[ID](#)43}, H.V. Cliff ^{[ID](#)50}, J. Closier ^{[ID](#)43}, J.L. Cobbledick ^{[ID](#)57}, V. Coco ^{[ID](#)43}, J. Cogan ^{[ID](#)10}, E. Cogneras ^{[ID](#)9}, L. Cojocariu ^{[ID](#)37}, P. Collins ^{[ID](#)43}, T. Colombo ^{[ID](#)43}, L. Congedo ^{[ID](#)19}, A. Contu ^{[ID](#)27}, N. Cooke ^{[ID](#)48}, I. Corredoira ^{[ID](#)41}, G. Corti ^{[ID](#)43}, B. Couturier ^{[ID](#)43}, D.C. Craik ^{[ID](#)45}, M. Cruz Torres ^{[ID](#)1,f}, R. Currie ^{[ID](#)53}, C.L. Da Silva ^{[ID](#)62}, S. Dadabaev ^{[ID](#)38}, L. Dai ^{[ID](#)66}, X. Dai ^{[ID](#)5}, E. Dall’Occo ^{[ID](#)15}, J. Dalseno ^{[ID](#)41}, C. D’Ambrosio ^{[ID](#)43}, J. Daniel ^{[ID](#)9}, A. Danilina ^{[ID](#)38}, P. d’Argent ^{[ID](#)19}, J.E. Davies ^{[ID](#)57}, A. Davis ^{[ID](#)57},

O. De Aguiar Francisco ⁵⁷, J. de Boer ⁴³, K. De Bruyn ⁷⁴, S. De Capua ⁵⁷, M. De Cian ⁴⁴,
U. De Freitas Carneiro Da Graca ¹, E. De Lucia ²³, J.M. De Miranda ¹, L. De Paula ²,
M. De Serio ^{19,g}, D. De Simone ⁴⁵, P. De Simone ²³, F. De Vellis ¹⁵, J.A. de Vries ⁷⁵,
C.T. Dean ⁶², F. Debernardis ^{19,g}, D. Decamp ⁸, V. Dedu ¹⁰, L. Del Buono ¹³,
B. Delaney ⁵⁹, H.-P. Dembinski ¹⁵, V. Denysenko ⁴⁵, O. Deschamps ⁹, F. Dettori ^{27,i},
B. Dey ⁷², P. Di Nezza ²³, I. Diachkov ³⁸, S. Didenko ³⁸, L. Dieste Maronas ⁴¹, S. Ding ⁶³,
V. Dobishuk ⁴⁷, A. Dolmatov ³⁸, C. Dong ³, A.M. Donohoe ¹⁸, F. Dordei ²⁷,
A.C. dos Reis ¹, L. Douglas ⁵⁴, A.G. Downes ⁸, P. Duda ⁷⁶, M.W. Dudek ³⁵, L. Dufour ⁴³,
V. Duk ⁷³, P. Durante ⁴³, M.M. Duras ⁷⁶, J.M. Durham ⁶², D. Dutta ⁵⁷, A. Dziurda ³⁵,
A. Dzyuba ³⁸, S. Easo ⁵², U. Egede ⁶⁴, A. Egorychev ³⁸, V. Egorychev ³⁸, C. Eirea Orro ⁴¹,
S. Eisenhardt ⁵³, E. Ejopu ⁵⁷, S. Ek-In ⁴⁴, L. Eklund ⁷⁷, M. Elashri ⁶⁰, J. Ellbracht ¹⁵,
S. Ely ⁵⁶, A. Ene ³⁷, E. Eppele ⁶⁰, S. Escher ¹⁴, J. Eschle ⁴⁵, S. Esen ⁴⁵, T. Evans ⁵⁷,
F. Fabiano ^{27,i}, L.N. Falcao ¹, Y. Fan ⁶, B. Fang ^{11,69}, L. Fantini ^{73,p}, M. Faria ⁴⁴,
S. Farry ⁵⁵, D. Fazzini ^{26,n}, L. Felkowski ⁷⁶, M. Feo ⁴³, M. Fernandez Gomez ⁴¹,
A.D. Fernez ⁶¹, F. Ferrari ²⁰, L. Ferreira Lopes ⁴⁴, F. Ferreira Rodrigues ²,
S. Ferreres Sole ³², M. Ferrillo ⁴⁵, M. Ferro-Luzzi ⁴³, S. Filippov ³⁸, R.A. Fini ¹⁹,
M. Fiorini ^{21,j}, M. Firlej ³⁴, K.M. Fischer ⁵⁸, D.S. Fitzgerald ⁷⁸, C. Fitzpatrick ⁵⁷,
T. Fiutowski ³⁴, F. Fleuret ¹², M. Fontana ²⁰, F. Fontanelli ^{24,l}, R. Forty ⁴³,
D. Foulds-Holt ⁵⁰, V. Franco Lima ⁵⁵, M. Franco Sevilla ⁶¹, M. Frank ⁴³, E. Franzoso ^{21,j},
G. Frau ¹⁷, C. Frei ⁴³, D.A. Friday ⁵⁷, L. Frontini ^{25,m}, J. Fu ⁶, Q. Fuehring ¹⁵,
T. Fulghesu ¹³, E. Gabriel ³², G. Galati ^{19,g}, M.D. Galati ³², A. Gallas Torreira ⁴¹,
D. Galli ^{20,h}, S. Gambetta ^{53,43}, M. Gandelman ², P. Gandini ²⁵, H. Gao ⁶, R. Gao ⁵⁸,
Y. Gao ⁷, Y. Gao ⁵, M. Garau ^{27,i}, L.M. Garcia Martin ⁵¹, P. Garcia Moreno ⁴⁰,
J. García Pardiñas ⁴³, B. Garcia Plana ⁴¹, F.A. Garcia Rosales ¹², L. Garrido ⁴⁰,
C. Gaspar ⁴³, R.E. Geertsema ³², D. Gerick ¹⁷, L.L. Gerken ¹⁵, E. Gersabeck ⁵⁷,
M. Gersabeck ⁵⁷, T. Gershon ⁵¹, L. Giambastiani ²⁸, V. Gibson ⁵⁰, H.K. Gienza ³⁶,
A.L. Gilman ⁵⁸, M. Giovannetti ²³, A. Gioventù ⁴¹, P. Gironella Gironell ⁴⁰,
C. Giugliano ^{21,j}, M.A. Giza ³⁵, K. Gizdov ⁵³, E.L. Gkougkousis ⁴³, V.V. Gligorov ^{13,43},
C. Göbel ⁶⁵, E. Golobardes ³⁹, D. Golubkov ³⁸, A. Golutvin ^{56,38}, A. Gomes ^{1,2,b,a,†},
S. Gomez Fernandez ⁴⁰, F. Goncalves Abrantes ⁵⁸, M. Goncerz ³⁵, G. Gong ³,
I.V. Gorelov ³⁸, C. Gotti ²⁶, J.P. Grabowski ⁷¹, T. Grammatico ¹³,
L.A. Granado Cardoso ⁴³, E. Graugés ⁴⁰, E. Graverini ⁴⁴, G. Graziani ¹⁰, A.T. Grecu ³⁷,
L.M. Greeven ³², N.A. Grieser ⁶⁰, L. Grillo ⁵⁴, S. Gromov ³⁸, C. Gu ³, M. Guarise ^{21,j},
M. Guittiere ¹¹, P.A. Günther ¹⁷, A.-K. Guseinov ³⁸, E. Gushchin ³⁸, A. Guth ¹⁴,
Y. Guz ^{5,38,43}, T. Gys ⁴³, T. Hadavizadeh ⁶⁴, C. Hadjivasiliou ⁶¹, G. Haefeli ⁴⁴,
C. Haen ⁴³, J. Haimberger ⁴³, S.C. Haines ⁵⁰, T. Halewood-leagas ⁵⁵, M.M. Halvorsen ⁴³,
P.M. Hamilton ⁶¹, J. Hammerich ⁵⁵, Q. Han ⁷, X. Han ¹⁷, S. Hansmann-Menzemer ¹⁷,
L. Hao ⁶, N. Harnew ⁵⁸, T. Harrison ⁵⁵, C. Hasse ⁴³, M. Hatch ⁴³, J. He ^{6,d},
K. Heijhoff ³², F. Hemmer ⁴³, C. Henderson ⁶⁰, R.D.L. Henderson ^{64,51},
A.M. Hennequin ⁵⁹, K. Hennessy ⁵⁵, L. Henry ⁴³, J. Herd ⁵⁶, J. Heuel ¹⁴, A. Hicheur ²,
D. Hill ⁴⁴, M. Hilton ⁵⁷, S.E. Hollitt ¹⁵, J. Horswill ⁵⁷, R. Hou ⁷, Y. Hou ⁸, J. Hu ¹⁷,
J. Hu ⁶⁷, W. Hu ⁵, X. Hu ³, W. Huang ⁶, X. Huang ⁶⁹, W. Hulsbergen ³², R.J. Hunter ⁵¹,
M. Hushchyn ³⁸, D. Hutchcroft ⁵⁵, P. Ibis ¹⁵, M. Idzik ³⁴, D. Ilin ³⁸, P. Ilten ⁶⁰,

A. Inglessi ^{ib38}, A. Iniukhin ^{ib38}, A. Ishteev ^{ib38}, K. Ivshin ^{ib38}, R. Jacobsson ^{ib43}, H. Jage ^{ib14}, S.J. Jaimes Elles ^{ib42}, S. Jakobsen ^{ib43}, E. Jans ^{ib32}, B.K. Jashal ^{ib42}, A. Jawahery ^{ib61}, V. Jevtic ^{ib15}, E. Jiang ^{ib61}, X. Jiang ^{ib4,6}, Y. Jiang ^{ib6}, M. John ^{ib58}, D. Johnson ^{ib59}, C.R. Jones ^{ib50}, T.P. Jones ^{ib51}, S. Joshi ^{ib36}, B. Jost ^{ib43}, N. Jurik ^{ib43}, I. Juszcak ^{ib35}, S. Kandybei ^{ib46}, Y. Kang ^{ib3}, M. Karacson ^{ib43}, D. Karpenkov ^{ib38}, M. Karpov ^{ib38}, J.W. Kautz ^{ib60}, F. Keizer ^{ib43}, D.M. Keller ^{ib63}, M. Kenzie ^{ib51}, T. Ketel ^{ib32}, B. Khanji ^{ib63}, A. Kharisova ^{ib38}, S. Kholodenko ^{ib38}, G. Khreich ^{ib11}, T. Kirn ^{ib14}, V.S. Kirsebom ^{ib44}, O. Kitouni ^{ib59}, S. Klaver ^{ib33}, N. Kleijne ^{ib29,q}, K. Klimaszewski ^{ib36}, M.R. Kmiec ^{ib36}, S. Koliiev ^{ib47}, L. Kolk ^{ib15}, A. Kondybayeva ^{ib38}, A. Konoplyannikov ^{ib38}, P. Kopciewicz ^{ib34}, R. Kopečna ^{ib17}, P. Koppenburg ^{ib32}, M. Korolev ^{ib38}, I. Kostiuk ^{ib32}, O. Kot ^{ib47}, S. Kotriakhova ^{ib}, A. Kozachuk ^{ib38}, P. Kravchenko ^{ib38}, L. Kravchuk ^{ib38}, M. Kreps ^{ib51}, S. Kretzschmar ^{ib14}, P. Krokovny ^{ib38}, W. Krupa ^{ib34}, W. Krzemien ^{ib36}, J. Kubat ^{ib17}, S. Kubis ^{ib76}, W. Kucewicz ^{ib35}, M. Kucharczyk ^{ib35}, V. Kudryavtsev ^{ib38}, E. Kulikova ^{ib38}, A. Kupsc ^{ib77}, D. Lacarrere ^{ib43}, G. Lafferty ^{ib57}, A. Lai ^{ib27}, A. Lampis ^{ib27,i}, D. Lancierini ^{ib45}, C. Landesa Gomez ^{ib41}, J.J. Lane ^{ib57}, R. Lane ^{ib49}, C. Langenbruch ^{ib14}, J. Langer ^{ib15}, O. Lantwin ^{ib38}, T. Latham ^{ib51}, F. Lazzari ^{ib29,r}, C. Lazzeroni ^{ib48}, R. Le Gac ^{ib10}, S.H. Lee ^{ib78}, R. Lefèvre ^{ib9}, A. Leflat ^{ib38}, S. Legotin ^{ib38}, O. Leroy ^{ib10}, T. Lesiak ^{ib35}, B. Leverington ^{ib17}, A. Li ^{ib3}, H. Li ^{ib67}, K. Li ^{ib7}, P. Li ^{ib43}, P.-R. Li ^{ib68}, S. Li ^{ib7}, T. Li ^{ib4}, T. Li ^{ib67}, Y. Li ^{ib4}, Z. Li ^{ib63}, X. Liang ^{ib63}, C. Lin ^{ib6}, T. Lin ^{ib52}, R. Lindner ^{ib43}, V. Lisovskyi ^{ib15}, R. Litvinov ^{ib27,i}, G. Liu ^{ib67}, H. Liu ^{ib6}, K. Liu ^{ib68}, Q. Liu ^{ib6}, S. Liu ^{ib4,6}, A. Lobo Salvia ^{ib40}, A. Loi ^{ib27}, R. Lollini ^{ib73}, J. Lomba Castro ^{ib41}, I. Longstaff ^{ib54}, J.H. Lopes ^{ib2}, A. Lopez Huertas ^{ib40}, S. López Soliño ^{ib41}, G.H. Lovell ^{ib50}, Y. Lu ^{ib4,c}, C. Lucarelli ^{ib22,k}, D. Lucchesi ^{ib28,o}, S. Luchuk ^{ib38}, M. Lucio Martinez ^{ib75}, V. Lukashenko ^{ib32,47}, Y. Luo ^{ib3}, A. Lupato ^{ib57}, E. Luppi ^{ib21,j}, A. Lusiani ^{ib29,q}, K. Lynch ^{ib18}, X.-R. Lyu ^{ib6}, R. Ma ^{ib6}, S. Maccolini ^{ib15}, F. Machefert ^{ib11}, F. Maciuc ^{ib37}, I. Mackay ^{ib58}, V. Macko ^{ib44}, L.R. Madhan Mohan ^{ib50}, A. Maevskiy ^{ib38}, D. Maisuzenko ^{ib38}, M.W. Majewski ^{ib34}, J.J. Malczewski ^{ib35}, S. Malde ^{ib58}, B. Malecki ^{ib35,43}, A. Malinin ^{ib38}, T. Maltsev ^{ib38}, G. Manca ^{ib27,i}, G. Mancinelli ^{ib10}, C. Mancuso ^{ib11,25,m}, R. Manera Escalero ^{ib40}, D. Manuzzi ^{ib20}, C.A. Manzari ^{ib45}, D. Marangotto ^{ib25,m}, J.F. Marchand ^{ib8}, U. Marconi ^{ib20}, S. Mariani ^{ib43}, C. Marin Benito ^{ib40}, J. Marks ^{ib17}, A.M. Marshall ^{ib49}, P.J. Marshall ^{ib55}, G. Martelli ^{ib73,p}, G. Martellotti ^{ib30}, L. Martinazzoli ^{ib43,n}, M. Martinelli ^{ib26,n}, D. Martinez Santos ^{ib41}, F. Martinez Vidal ^{ib42}, A. Massafferri ^{ib1}, M. Materok ^{ib14}, R. Matev ^{ib43}, A. Mathad ^{ib45}, V. Matiunin ^{ib38}, C. Matteuzzi ^{ib26}, K.R. Mattioli ^{ib12}, A. Mauri ^{ib56}, E. Maurice ^{ib12}, J. Mauricio ^{ib40}, M. Mazurek ^{ib43}, M. McCann ^{ib56}, L. Mcconnell ^{ib18}, T.H. McGrath ^{ib57}, N.T. McHugh ^{ib54}, A. McNab ^{ib57}, R. McNulty ^{ib18}, B. Meadows ^{ib60}, G. Meier ^{ib15}, D. Melnychuk ^{ib36}, S. Meloni ^{ib26,n}, M. Merk ^{ib32,75}, A. Merli ^{ib25,m}, L. Meyer Garcia ^{ib2}, D. Miao ^{ib4,6}, H. Miao ^{ib6}, M. Mikhasenko ^{ib71,e}, D.A. Milanes ^{ib70}, E. Millard ^{ib51}, M. Milovanovic ^{ib43}, M.-N. Minard ^{ib8,†}, A. Minotti ^{ib26,n}, E. Minucci ^{ib63}, T. Miralles ^{ib9}, S.E. Mitchell ^{ib53}, B. Mitreska ^{ib15}, D.S. Mitzel ^{ib15}, A. Modak ^{ib52}, A. Mödden ^{ib15}, R.A. Mohammed ^{ib58}, R.D. Moise ^{ib14}, S. Mokhnenko ^{ib38}, T. Mombächer ^{ib41}, M. Monk ^{ib51,64}, I.A. Monroy ^{ib70}, S. Monteil ^{ib9}, G. Morello ^{ib23}, M.J. Morello ^{ib29,q}, M.P. Morgenthaler ^{ib17}, J. Moron ^{ib34}, A.B. Morris ^{ib43}, A.G. Morris ^{ib10}, R. Mountain ^{ib63}, H. Mu ^{ib3}, E. Muhammad ^{ib51}, F. Muheim ^{ib53}, M. Mulder ^{ib74}, K. Müller ^{ib45}, C.H. Murphy ^{ib58}, D. Murray ^{ib57}, R. Murta ^{ib56}, P. Muzzetto ^{ib27,i}, P. Naik ^{ib49}, T. Nakada ^{ib44}, R. Nandakumar ^{ib52}, T. Nanut ^{ib43}, I. Nasteva ^{ib2},

M. Needham ⁵³, N. Neri ^{25,m}, S. Neubert ⁷¹, N. Neufeld ⁴³, P. Neustroev ³⁸, R. Newcombe ⁵⁶, J. Nicolini ^{15,11}, D. Nicotra ⁷⁵, E.M. Niel ⁴⁴, S. Nieswand ¹⁴, N. Nikitin ³⁸, N.S. Nolte ⁵⁹, C. Normand ^{8,i,27}, J. Novoa Fernandez ⁴¹, G. Nowak ⁶⁰, C. Nunez ⁷⁸, A. Oblakowska-Mucha ³⁴, V. Obraztsov ³⁸, T. Oeser ¹⁴, S. Okamura ^{21,j}, R. Oldeman ^{27,i}, F. Oliva ⁵³, C.J.G. Onderwater ⁷⁴, R.H. O’Neil ⁵³, J.M. Otalora Goicochea ², T. Ovsianikova ³⁸, P. Owen ⁴⁵, A. Oyanguren ⁴², O. Ozcelik ⁵³, K.O. Padeken ⁷¹, B. Pagare ⁵¹, P.R. Pais ⁴³, T. Pajero ⁵⁸, A. Palano ¹⁹, M. Palutan ²³, G. Panshin ³⁸, L. Paolucci ⁵¹, A. Papanestis ⁵², M. Pappagallo ^{19,g}, L.L. Pappalardo ^{21,j}, C. Pappenheimer ⁶⁰, W. Parker ⁶¹, C. Parkes ^{57,43}, B. Passalacqua ^{21,j}, G. Passaleva ²², A. Pastore ¹⁹, M. Patel ⁵⁶, C. Patrignani ^{20,h}, C.J. Pawley ⁷⁵, A. Pellegrino ³², M. Pepe Altarelli ⁴³, S. Perazzini ²⁰, D. Pereima ³⁸, A. Pereiro Castro ⁴¹, P. Perret ⁹, K. Petridis ⁴⁹, A. Petrolini ^{24,l}, S. Petrucci ⁵³, M. Petruzzo ²⁵, H. Pham ⁶³, A. Philippov ³⁸, R. Piandani ⁶, L. Pica ^{29,q}, M. Piccini ⁷³, B. Pietrzyk ⁸, G. Pietrzyk ¹¹, M. Pili ⁵⁸, D. Pinci ³⁰, F. Pisani ⁴³, M. Pizzichemi ^{26,n,43}, V. Placinta ³⁷, J. Plews ⁴⁸, M. Plo Casasus ⁴¹, F. Polci ^{13,43}, M. Poli Lener ²³, A. Poluektov ¹⁰, N. Polukhina ³⁸, I. Polyakov ⁴³, E. Polycarpo ², S. Ponce ⁴³, D. Popov ^{6,43}, S. Poslavskii ³⁸, K. Prasanth ³⁵, L. Promberger ¹⁷, C. Prouve ⁴¹, V. Pugatch ⁴⁷, V. Puill ¹¹, G. Punzi ^{29,r}, H.R. Qi ³, W. Qian ⁶, N. Qin ³, S. Qu ³, R. Quagliani ⁴⁴, N.V. Raab ¹⁸, B. Rachwal ³⁴, J.H. Rademacker ⁴⁹, R. Rajagopalan ⁶³, M. Rama ²⁹, M. Ramos Pernas ⁵¹, M.S. Rangel ², F. Ratnikov ³⁸, G. Raven ³³, M. Rebollo De Miguel ⁴², F. Redi ⁴³, J. Reich ⁴⁹, F. Reiss ⁵⁷, C. Remon Alepuz ⁴², Z. Ren ³, P.K. Resmi ⁵⁸, R. Ribatti ^{29,q}, A.M. Ricci ²⁷, S. Ricciardi ⁵², K. Richardson ⁵⁹, M. Richardson-Slipper ⁵³, K. Rinnert ⁵⁵, P. Robbe ¹¹, G. Robertson ⁵³, E. Rodrigues ^{55,43}, E. Rodriguez Fernandez ⁴¹, J.A. Rodriguez Lopez ⁷⁰, E. Rodriguez Rodriguez ⁴¹, D.L. Rolf ⁴³, A. Rollings ⁵⁸, P. Roloff ⁴³, V. Romanovskiy ³⁸, M. Romero Lamas ⁴¹, A. Romero Vidal ⁴¹, M. Rotondo ²³, M.S. Rudolph ⁶³, T. Ruf ⁴³, R.A. Ruiz Fernandez ⁴¹, J. Ruiz Vidal ⁴², A. Ryzhikov ³⁸, J. Ryzka ³⁴, J.J. Saborido Silva ⁴¹, N. Sagidova ³⁸, N. Sahoo ⁴⁸, B. Saitta ^{27,i}, M. Salomoni ⁴³, C. Sanchez Gras ³², I. Sanderswood ⁴², R. Santacesaria ³⁰, C. Santamarina Rios ⁴¹, M. Santimaria ²³, L. Santoro ¹, E. Santovetti ^{31,t}, D. Saranin ³⁸, G. Sarpis ⁵³, M. Sarpis ⁷¹, A. Sarti ³⁰, C. Satriano ^{30,s}, A. Satta ³¹, M. Saur ⁵, D. Savrina ³⁸, H. Sazak ⁹, L.G. Scantlebury Smead ⁵⁸, A. Scarabotto ¹³, S. Schael ¹⁴, S. Scherl ⁵⁵, A.M. Schertz ⁷², M. Schiller ⁵⁴, H. Schindler ⁴³, M. Schmelling ¹⁶, B. Schmidt ⁴³, S. Schmitt ¹⁴, O. Schneider ⁴⁴, A. Schopper ⁴³, M. Schubiger ³², N. Schulte ¹⁵, S. Schulte ⁴⁴, M.H. Schune ¹¹, R. Schwemmer ⁴³, B. Sciascia ²³, A. Sciuccati ⁴³, S. Sellam ⁴¹, A. Semennikov ³⁸, M. Senghi Soares ³³, A. Sergi ^{24,l}, N. Serra ⁴⁵, L. Sestini ²⁸, A. Seuthe ¹⁵, Y. Shang ⁵, D.M. Shangase ⁷⁸, M. Shapkin ³⁸, I. Shchemerov ³⁸, L. Shchutska ⁴⁴, T. Shears ⁵⁵, L. Shekhtman ³⁸, Z. Shen ⁵, S. Sheng ^{4,6}, V. Shevchenko ³⁸, B. Shi ⁶, E.B. Shields ^{26,n}, Y. Shimizu ¹¹, E. Shmanin ³⁸, R. Shorkin ³⁸, J.D. Shupperd ⁶³, B.G. Siddi ^{21,j}, R. Silva Coutinho ⁶³, G. Simi ²⁸, S. Simone ^{19,g}, M. Singla ⁶⁴, N. Skidmore ⁵⁷, R. Skuza ¹⁷, T. Skwarnicki ⁶³, M.W. Slater ⁴⁸, J.C. Smallwood ⁵⁸, J.G. Smeaton ⁵⁰, E. Smith ⁴⁵, K. Smith ⁶², M. Smith ⁵⁶, A. Snoch ³², L. Soares Lavra ⁹, M.D. Sokoloff ⁶⁰, F.J.P. Soler ⁵⁴, A. Solomin ^{38,49}, A. Solovev ³⁸, I. Solovyev ³⁸, R. Song ⁶⁴, F.L. Souza De Almeida ²,

B. Souza De Paula ², E. Spadaro Norella ^{25,m}, E. Spedicato ²⁰, J.G. Speer ¹⁵,
 E. Spiridenkov ³⁸, P. Spradlin ⁵⁴, V. Sriskaran ⁴³, F. Stagni ⁴³, M. Stahl ⁴³, S. Stahl ⁴³,
 S. Stanislaus ⁵⁸, E.N. Stein ⁴³, O. Steinkamp ⁴⁵, O. Stenyakin ³⁸, H. Stevens ¹⁵,
 D. Strekalina ³⁸, Y. Su ⁶, F. Suljik ⁵⁸, J. Sun ²⁷, L. Sun ⁶⁹, Y. Sun ⁶¹, P.N. Swallow ⁴⁸,
 K. Swientek ³⁴, A. Szabelski ³⁶, T. Szumlak ³⁴, M. Szymanski ⁴³, Y. Tan ³, S. Taneja ⁵⁷,
 M.D. Tat ⁵⁸, A. Terentev ⁴⁵, F. Teubert ⁴³, E. Thomas ⁴³, D.J.D. Thompson ⁴⁸,
 H. Tilquin ⁵⁶, V. Tisserand ⁹, S. T'Jampens ⁸, M. Tobin ⁴, L. Tomassetti ^{21,j},
 G. Tonani ^{25,m}, X. Tong ⁵, D. Torres Machado ¹, D.Y. Tou ³, C. Trippel ⁴⁴, G. Tuci ⁶,
 N. Tuning ³², A. Ukleja ³⁶, D.J. Unverzagt ¹⁷, A. Usachov ³³, A. Ustyuzhanin ³⁸,
 U. Uwer ¹⁷, V. Vagnoni ²⁰, A. Valassi ⁴³, G. Valenti ²⁰, N. Valls Canudas ³⁹,
 M. Van Dijk ⁴⁴, H. Van Hecke ⁶², E. van Herwijnen ⁵⁶, C.B. Van Hulse ^{41,v},
 M. van Veghel ³², R. Vazquez Gomez ⁴⁰, P. Vazquez Regueiro ⁴¹, C. Vázquez Sierra ⁴¹,
 S. Vecchi ²¹, J.J. Velthuis ⁴⁹, M. Veltri ^{22,u}, A. Venkateswaran ⁴⁴, M. Veronesi ³²,
 M. Vesterinen ⁵¹, D. Vieira ⁶⁰, M. Vieites Diaz ⁴⁴, X. Vilasis-Cardona ³⁹,
 E. Vilella Figueras ⁵⁵, A. Villa ²⁰, P. Vincent ¹³, F.C. Volle ¹¹, D. vom Bruch ¹⁰,
 V. Vorobyev ³⁸, N. Voropaev ³⁸, K. Vos ⁷⁵, C. Vrahas ⁵³, J. Walsh ²⁹, E.J. Walton ⁶⁴,
 G. Wan ⁵, C. Wang ¹⁷, G. Wang ⁷, J. Wang ⁵, J. Wang ⁴, J. Wang ³, J. Wang ⁶⁹,
 M. Wang ²⁵, R. Wang ⁴⁹, X. Wang ⁶⁷, Y. Wang ⁷, Z. Wang ⁴⁵, Z. Wang ³, Z. Wang ⁶,
 J.A. Ward ^{51,64}, N.K. Watson ⁴⁸, D. Websdale ⁵⁶, Y. Wei ⁵, B.D.C. Westhenry ⁴⁹,
 D.J. White ⁵⁷, M. Whitehead ⁵⁴, A.R. Wiederhold ⁵¹, D. Wiedner ¹⁵, G. Wilkinson ⁵⁸,
 M.K. Wilkinson ⁶⁰, I. Williams ⁵⁰, M. Williams ⁵⁹, M.R.J. Williams ⁵³, R. Williams ⁵⁰,
 F.F. Wilson ⁵², W. Wislicki ³⁶, M. Witek ³⁵, L. Witola ¹⁷, C.P. Wong ⁶², G. Wormser ¹¹,
 S.A. Wotton ⁵⁰, H. Wu ⁶³, J. Wu ⁷, K. Wyllie ⁴³, Z. Xiang ⁶, Y. Xie ⁷, A. Xu ⁵,
 J. Xu ⁶, L. Xu ³, L. Xu ³, M. Xu ⁵¹, Q. Xu ⁶, Z. Xu ⁹, Z. Xu ⁶, D. Yang ³, S. Yang ⁶,
 X. Yang ⁵, Y. Yang ⁶, Z. Yang ⁵, Z. Yang ⁶¹, L.E. Yeomans ⁵⁵, V. Yeroshenko ¹¹,
 H. Yeung ⁵⁷, H. Yin ⁷, J. Yu ⁶⁶, X. Yuan ⁶³, E. Zaffaroni ⁴⁴, M. Zavertyaev ¹⁶,
 M. Zdybal ³⁵, M. Zeng ³, C. Zhang ⁵, D. Zhang ⁷, J. Zhang ⁶, L. Zhang ³, S. Zhang ⁶⁶,
 S. Zhang ⁵, Y. Zhang ⁵, Y. Zhang ⁵⁸, Y. Zhao ¹⁷, A. Zharkova ³⁸, A. Zhelezov ¹⁷,
 Y. Zheng ⁶, T. Zhou ⁵, X. Zhou ⁷, Y. Zhou ⁶, V. Zhovkovska ¹¹, X. Zhu ³, X. Zhu ⁷,
 Z. Zhu ⁶, V. Zhukov ^{14,38}, J. Zhuo ⁴², Q. Zou ^{4,6}, S. Zucchelli ^{20,h}, D. Zuliani ²⁸,
 G. Zunica ⁵⁷

¹ Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

² Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³ Center for High Energy Physics, Tsinghua University, Beijing, China

⁴ Institute Of High Energy Physics (IHEP), Beijing, China

⁵ School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

⁶ University of Chinese Academy of Sciences, Beijing, China

⁷ Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China

⁸ Université Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

⁹ Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

¹⁰ Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

¹¹ Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France

¹² Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

- ¹³ LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France
- ¹⁴ I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany
- ¹⁵ Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany
- ¹⁶ Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany
- ¹⁷ Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- ¹⁸ School of Physics, University College Dublin, Dublin, Ireland
- ¹⁹ INFN Sezione di Bari, Bari, Italy
- ²⁰ INFN Sezione di Bologna, Bologna, Italy
- ²¹ INFN Sezione di Ferrara, Ferrara, Italy
- ²² INFN Sezione di Firenze, Firenze, Italy
- ²³ INFN Laboratori Nazionali di Frascati, Frascati, Italy
- ²⁴ INFN Sezione di Genova, Genova, Italy
- ²⁵ INFN Sezione di Milano, Milano, Italy
- ²⁶ INFN Sezione di Milano-Bicocca, Milano, Italy
- ²⁷ INFN Sezione di Cagliari, Monserrato, Italy
- ²⁸ Università degli Studi di Padova, Università e INFN, Padova, Padova, Italy
- ²⁹ INFN Sezione di Pisa, Pisa, Italy
- ³⁰ INFN Sezione di Roma La Sapienza, Roma, Italy
- ³¹ INFN Sezione di Roma Tor Vergata, Roma, Italy
- ³² Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands
- ³³ Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands
- ³⁴ AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland
- ³⁵ Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland
- ³⁶ National Center for Nuclear Research (NCBJ), Warsaw, Poland
- ³⁷ Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania
- ³⁸ Affiliated with an institute covered by a cooperation agreement with CERN
- ³⁹ DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain
- ⁴⁰ ICCUB, Universitat de Barcelona, Barcelona, Spain
- ⁴¹ Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain
- ⁴² Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain
- ⁴³ European Organization for Nuclear Research (CERN), Geneva, Switzerland
- ⁴⁴ Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland
- ⁴⁵ Physik-Institut, Universität Zürich, Zürich, Switzerland
- ⁴⁶ NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine
- ⁴⁷ Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine
- ⁴⁸ University of Birmingham, Birmingham, United Kingdom
- ⁴⁹ H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom
- ⁵⁰ Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom
- ⁵¹ Department of Physics, University of Warwick, Coventry, United Kingdom
- ⁵² STFC Rutherford Appleton Laboratory, Didcot, United Kingdom
- ⁵³ School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom
- ⁵⁴ School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom
- ⁵⁵ Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom
- ⁵⁶ Imperial College London, London, United Kingdom
- ⁵⁷ Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom
- ⁵⁸ Department of Physics, University of Oxford, Oxford, United Kingdom
- ⁵⁹ Massachusetts Institute of Technology, Cambridge, MA, United States
- ⁶⁰ University of Cincinnati, Cincinnati, OH, United States
- ⁶¹ University of Maryland, College Park, MD, United States

- ⁶² Los Alamos National Laboratory (LANL), Los Alamos, NM, United States
- ⁶³ Syracuse University, Syracuse, NY, United States
- ⁶⁴ School of Physics and Astronomy, Monash University, Melbourne, Australia, associated to ⁵¹
- ⁶⁵ Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²
- ⁶⁶ Physics and Micro Electronic College, Hunan University, Changsha City, China, associated to ⁷
- ⁶⁷ Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Institute of Quantum Matter, South China Normal University, Guangzhou, China, associated to ³
- ⁶⁸ Lanzhou University, Lanzhou, China, associated to ⁴
- ⁶⁹ School of Physics and Technology, Wuhan University, Wuhan, China, associated to ³
- ⁷⁰ Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to ¹³
- ⁷¹ Universität Bonn - Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany, associated to ¹⁷
- ⁷² Eotvos Lorand University, Budapest, Hungary, associated to ⁴³
- ⁷³ INFN Sezione di Perugia, Perugia, Italy, associated to ²¹
- ⁷⁴ Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to ³²
- ⁷⁵ Universiteit Maastricht, Maastricht, Netherlands, associated to ³²
- ⁷⁶ Tadeusz Kosciuszko Cracow University of Technology, Cracow, Poland, associated to ³⁵
- ⁷⁷ Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden, associated to ⁵⁴
- ⁷⁸ University of Michigan, Ann Arbor, MI, United States, associated to ⁶³

^a Universidade de Brasília, Brasília, Brazil

^b Universidade Federal do Triângulo Mineiro (UFMT), Uberaba-MG, Brazil

^c Central South U., Changsha, China

^d Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China

^e Excellence Cluster ORIGINS, Munich, Germany

^f Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras

^g Università di Bari, Bari, Italy

^h Università di Bologna, Bologna, Italy

ⁱ Università di Cagliari, Cagliari, Italy

^j Università di Ferrara, Ferrara, Italy

^k Università di Firenze, Firenze, Italy

^l Università di Genova, Genova, Italy

^m Università degli Studi di Milano, Milano, Italy

ⁿ Università di Milano Bicocca, Milano, Italy

^o Università di Padova, Padova, Italy

^p Università di Perugia, Perugia, Italy

^q Scuola Normale Superiore, Pisa, Italy

^r Università di Pisa, Pisa, Italy

^s Università della Basilicata, Potenza, Italy

^t Università di Roma Tor Vergata, Roma, Italy

^u Università di Urbino, Urbino, Italy

^v Universidad de Alcalá, Alcalá de Henares, Spain

[†] Deceased